

Western Australian Immunisation Schedule

Childhood

| Age | Antigen | Vaccine brand(s) | Important information |
|------------------------------------|---|------------------------------------|--|
| Birth to <7 days | Hepatitis B | H-B-Vax® II Paed or Engerix®B Paed | |
| 6-8 Weeks | DTPa-hepB-IPV-Hib | Infanrix® hexa | |
| | 13vPCV | Prevenar® 13 | |
| | Rotavirus | Rotarix® | Oral dose 1: 6–14 weeks of age |
| Aboriginal | MenACWY | Nimenrix® | |
| Aboriginal and/or MRC | MenB | Bexsero® | Prophylactic paracetamol recommended prior to dose |
| 4 Months | DTPa-hepB-IPV-Hib | Infanrix® hexa | |
| | 13vPCV | Prevenar® 13 | |
| | Rotavirus | Rotarix® | Oral dose 2: 10–24 weeks of age |
| Aboriginal | MenACWY | Nimenrix® | |
| Aboriginal and/or MRC | MenB | Bexsero® | Prophylactic paracetamol recommended prior to dose |
| 6 Months | DTPa-hepB-IPV-Hib | Infanrix® hexa | |
| Aboriginal and/or MRC | 13vPCV | Prevenar® 13 | |
| MRC | MenB | Bexsero® | Prophylactic paracetamol recommended prior to dose |
| 12 Months | MMR | MMRII® or Priorix® | |
| | 13vPCV | Prevenar® 13 | |
| | MenACWY | Nimenrix® * | |
| Aboriginal and/or MRC | MenB | Bexsero® | Prophylactic paracetamol recommended prior to dose |
| MRC | Hepatitis B | H-B-Vax® II Paed or Engerix®B Paed | <32 weeks gestation or <2000g birth weight |
| All children > 12 months | Children diagnosed with MRC at >12 months of age require 1 dose of 13vPCV at diagnosis, followed by a first dose of 23vPPV, 2 to 12 months later or at 4 years of age, whichever is later. A second dose of 23vPPV is given 5 years later. See Australian Immunisation Handbook | | |
| MRC | | | |
| 18 Months | DTPa | Infanrix® or Tripacel® | |
| | Hib | ActHIB® | |
| | MMRV | Priorix-Tetra® or ProQuad® | Do not administer MMRV as dose 1 of MMR containing vaccine if <4 years of age |
| Aboriginal | Hepatitis A | Vaqa® Paed/Adol | First dose of the 2 dose schedule |
| <2 Years | Catch-up for MenB is available for Aboriginal children <2 years of age until 30 June 2023. | | |
| Aboriginal | | | |
| 4 Years | DTPa-IPV | Quadracel® or Infanrix®IPV | |
| Aboriginal | Hepatitis A | Vaqa® Paed/Adol | Not required if previously received 2 doses (first dose at age ≥12 months) at least 6 months apart |
| | 23vPPV | Pneumovax23® | First dose (if not previously received) Second dose at least 5 years later |

| Age | Antigen | Vaccine brand(s) | Important information |
|-------------------------------------|--|------------------------------|--|
| All adolescents and adults with MRC | Adolescents and adults diagnosed with MRC require 1 dose of 13vPCV at diagnosis, followed by a first dose of 23vPPV, 2 to 12 months later. A second dose is given at least 5 years later. People of all ages with specified MRC are eligible for MenACWY, MenB and Hib vaccines. See the ATAGI clinical advice for pneumococcal and meningococcal diseases as well as the Australian Immunisation Handbook for more information. | | |
| Year 7 | HPV | Gardasil®9 | 2 doses of Gardasil®9, 6 months apart |
| | dTpa | Boostrix® | |
| Year 10 | MenACWY | Nimenrix® | Catch-up for adolescents <20 years of age |
| Pregnant | dTpa | Boostrix® or Adacel® | Each pregnancy, ideally between 20–32 weeks |
| ≥20 Years Aboriginal | Hepatitis B | Engerix-B (adult) | For non-immune persons. See the Australian Immunisation Handbook for recommendations and timing of doses |
| ≥50 Years Aboriginal | 13vPCV 23vPPV | Prevenar® 13 Pneumovax23® | See the Australian Immunisation Handbook for for timing of doses |
| Born after 1965 | MMR | MMRII® or Priorix® | Give 1 or 2 doses as required |
| ≥70 Years | 13vPCV | Prevenar® 13 | |
| ≥70 to 79 Years | Herpes-zoster | Zostavax® | Catch-up program for 71–79 year olds until 31 Oct 2021 |

2020 Seasonal influenza

| Age | Vaccine brand(s) | Important information |
|---|--|--|
| 6 Months to <5 Years | Vaxigrip® Tetra, Fluarix® Tetra, FluQuadri® | If child <9 years, 2 doses required 1 month apart in the first year they receive influenza vaccine |
| ≥5 Years to Year 6 | FluQuadri® (WA State Program) | |
| 5 to <65 Years Aboriginal and/or MRC | Afluria® Quad, Fluarix® Tetra, FluQuadri®, Vaxigrip® Tetra | |
| Pregnant | Afluria® Quad, Fluarix® Tetra, FluQuadri®, Vaxigrip® Tetra | Each pregnancy, any trimester |
| ≥65 Years | Fluad® Quad | Use standard QIV if Fluad® Quad unavailable |
| WA Health staff | Fluarix® Tetra, FluQuadri® (WA State Program) | For ≥65 years use Fluad® Quad |

The term 'Aboriginal' is inclusive of all Aboriginal and Torres Strait Islander people.

MRC refers to medical risk conditions. Refer to the current online edition of The [Australian Immunisation Handbook](#) for all medical risk conditions.

Persons aged <20 years can catch-up scheduled vaccines, refer to [Australian Immunisation Handbook](#) for timing of doses.

Adult refugees and humanitarian entrants are eligible for dTpa, Hepatitis B, IPV, MMR and VZV.

* **MenACWY catch-up** for Aboriginal children <5 years of age until 31st January 2023.