

<input checked="" type="checkbox"/>	<u>CONSULTATION</u>	
<input type="checkbox"/>	<u>Person(s) present for consult:</u>	<input type="text"/>
<input type="checkbox"/>	<u>EMBRYO TRANSFER:</u>	
<input checked="" type="checkbox"/>		Number of embryos available for transfer: <input type="text"/>
<input checked="" type="checkbox"/>		Number of embryos to transfer: <input type="text"/>
<input checked="" type="checkbox"/>		Embryo Stage and Grade: <input type="text"/>
<input checked="" type="checkbox"/>		We discussed the risks of twins is approximately 30 - 40 percent and triplet
<input checked="" type="checkbox"/>		We discussed that the option of selective reduction carries a risk of caus
<input checked="" type="checkbox"/>		Any remaining viable embryos will be cultured and/or cryopreserved. All
<input type="checkbox"/>		Assisted hatching was discussed and will be performed.
<input checked="" type="checkbox"/>		Valium
	<input checked="" type="radio"/>	10-mg
	<input type="radio"/>	5-mg
	<input type="radio"/>	15-mg
	<input type="radio"/>	20-mg
<input type="checkbox"/>		Comment: <input type="text"/>
<input checked="" type="checkbox"/>		See ET report and signed consent for details.
<input type="checkbox"/>	<u>FET:</u>	
<input checked="" type="checkbox"/>		Number of embryos available for FET: <input type="text"/>
<input checked="" type="checkbox"/>		We discussed the risks of twins is approximately 20 - 25 percent and triplet
<input checked="" type="checkbox"/>		We discussed that the option of selective reduction carries a risk of caus
<input checked="" type="checkbox"/>		Assisted hatching was discussed and will be performed.
<input checked="" type="checkbox"/>		Valium
	<input checked="" type="radio"/>	10-mg
	<input type="radio"/>	5-mg
	<input type="radio"/>	15-mg
	<input type="radio"/>	20-mg
<input type="checkbox"/>		Comment: <input type="text"/>
<input checked="" type="checkbox"/>		See FET report and signed consent for details.
<input type="checkbox"/>	<u>OFFICE HYSTEROSCOPY:</u>	
<input checked="" type="checkbox"/>		We discussed the procedure for office hysteroscopy. The risks include bl
<input type="checkbox"/>		When CO2 is used, the risks also include CO2 embolus.
<input checked="" type="checkbox"/>		See signed consent and report.
<input checked="" type="checkbox"/>		Medications:
	<input checked="" type="checkbox"/>	doxycycline prophylaxis
	<input checked="" type="checkbox"/>	Toradol IM
	<input checked="" type="radio"/>	
	<input type="radio"/>	
	<input checked="" type="checkbox"/>	Valium
	<input checked="" type="radio"/>	
	<input type="radio"/>	
	<input type="radio"/>	
	<input type="radio"/>	
	<input type="checkbox"/>	Vicodin
	<input type="radio"/>	
	<input type="radio"/>	
	<input type="checkbox"/>	Ibuprofen
	<input type="checkbox"/>	Other medications:
<input type="checkbox"/>		Comment: <input type="text"/>
<input type="checkbox"/>	<u>Laparoscopy and Hysteroscopy</u>	

<input checked="" type="checkbox"/>	tubal dye studies	
<input type="checkbox"/>	and:	
<input checked="" type="checkbox"/>	We discussed the above procedure, alternatives and risks. The risks incl	
<input type="checkbox"/>	The risks of myomectomy also include, pelvic/tubal/intra-uterine adhesio	
<input checked="" type="checkbox"/>	All questions were answered.	
<input checked="" type="checkbox"/>	See H&P and separate signed consent.	
	<input type="radio"/>	Greater Sacramento Surger
	<input type="radio"/>	Mercy San Juan
	<input type="radio"/>	Sutter Roseville
<input checked="" type="checkbox"/>	Post-op medication prescription given to Pt. for	
	<input checked="" type="checkbox"/>	ibuprofen 800 mg po q 6 hr.
		<input checked="" type="checkbox"/>
		<input checked="" type="checkbox"/>
		<input type="checkbox"/>
		<input type="checkbox"/>
	<input checked="" type="checkbox"/>	Vicodin 1-2 tabs. q 6 hr., prn
		<input checked="" type="checkbox"/>
		<input checked="" type="checkbox"/>
		<input type="checkbox"/>
		<input type="checkbox"/>
	<input checked="" type="checkbox"/>	Phenergen 25 mg po q 6 hr,
		<input checked="" type="checkbox"/>
		<input checked="" type="checkbox"/>
	<input type="checkbox"/>	Other:
<input type="checkbox"/>	Comment:	

percent and triplets, or more, is less than 5 percent. There is a small risk of ectopic pregnancy.
es a risk of causing a miscarriage.
opreserved. All other non-viable eggs and embryos have been, or will be, discarded.

percent and triplets, or more, is less than 5 percent. There is a small risk of ectopic pregnancy.
es a risk of causing a miscarriage.

risks include but are not limited to infection, bleeding, uterine perforation, possibly requiring hospitalization or surgical correction

prophylaxis

30-mg

60-mg

10-mg

5-mg

15-mg

20-mg

1 tablet

2 tablets

cations:

s. The risks include, but are not limited to, the risk of anesthesia, infection, bleeding and damage to nearby organs. It may not be possible to avoid uterine adhesions, which may require a second procedure to correct. The need for Cesarean section, without labor. A less than on

San Francisco Surgery Center

Juan

ville

100 mg po q 6 hr.

Disp. No. 30

Refill x 1

Disp. No.

Refill x

100 mg tabs. q 6 hr., prn

Disp. No. 30

No Refill

Disp. No.

Refill x

25 mg po q 6 hr, prn

Disp. No 6

No Refill