

Product Category: **Similac®**

Bottles & Caps
Liquid Protein Fortifier
Similac Expert Care® 24 Cal With Iron
Similac Expert Care® Alimentum®
Similac Expert Care® for Diarrhea
Similac Expert Care® NeoSure®
Similac For Spit-Up®
Similac Go & Grow® Milk-Based Formula
Similac Go & Grow® Soy-Based Formula
Similac Sensitive®
Similac Total Comfort™
Similac® 10-pct Glucose Water
Similac® 5-pct Glucose Water
Similac® Advance®
Similac® Advance® Organic
Similac® For Supplementation
Similac® Human Milk Fortifier
Similac® Human Milk Fortifier Concentrated Liquid
Similac® Nipples
Similac® PM 60/40
Similac® Prenatal & Breastfeeding Dietary Supplement
Similac® Simply Smart™
Similac® Soy Isomil®
Similac® Sterilized Water
Similac® Volu-Feed®

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Product Information: Bottles & Caps

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Bottles & Caps

- Plastic (polypropylene) bottles designed for use in storing human milk. Standard (40-mm) plastic bottle caps fit on 2-fl-oz and 4-fl-oz human milk storage bottles, the 8-fl-oz plastic bottles and Volu-Feed® Nurser.
- Bottles are ready for use—do not require sterilization before first use.
- Dishwasher safe (top rack).
- DEHP- and Bisphenol A-free.



Availability

List Number	Item
00875	8-fl-oz Plastic Bottles / 144 ct
51950	4-fl-oz Plastic Bottles with Caps / 48 x 3 ct
54080	Bottle Caps / 250 ct
57752	2-fl-oz Plastic Bottles with Caps & Clips / 36 x 4 ct

Preparation

Cleanliness is important. Before preparing and filling bottles, wash your hands thoroughly.

Bottles DO NOT require sterilization before first use. AFTER FIRST USE, follow these instructions for preparation and use:

- Wash bottles and caps in hot, soapy water and rinse thoroughly; or wash in top rack of dishwasher.
- Your healthcare professional may recommend that you sterilize bottles before filling them with breast milk.
- Fill bottles with breast milk. Cap and refrigerate or freeze them promptly. If bottles are not filled immediately, replace caps until needed.

Rinse bottle and cap following use to prevent milk from sticking to bottle.

Filled bottles should be stored in refrigerator and used as directed by your healthcare professional. Breast milk should be used or frozen within 48 hours of expressing. Leave some empty space at the top of the bottle for milk to expand as it freezes. Label all bottles with name, date, and time. When ready to use frozen breast milk, swirl bottle under warm tap water to thaw. Feed thawed breast milk right away or refrigerate; do not refreeze.

WARNING: Never use a microwave oven to thaw or heat milk. Serious burns can result.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Product Information: Liquid Protein Fortifier

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Liquid Protein Fortifier

Extensively Hydrolyzed Protein

- For use with human milk or infant formula.
- The first and only commercially sterile, extensively hydrolyzed liquid protein designed for preterm infant feedings.
- Peptides and amino acids for easy digestion and absorption.
- Eliminates the need for powder mixing and meets ADA and CDC recommendations to reduce risk of contamination.^{1,2}
- Minimal impact on osmolality; 1 mL contributes ~12 mOsm/kg water when 100 mL feeding is made.

Safety Precautions

- Must be mixed with human milk or infant formula.
- This product is nutritionally incomplete.
- Do not attach nipple and ring.
- Enteral use only; not for IV use.
- **If signs of intolerance develop, slow feeding or discontinue.**

¹ Robbins ST, Beker LT. *Infant Feedings: Guidelines for Preparation of Formula and Breastmilk in Health Care Facilities*. American Dietetic Association, 2004.

² Centers for Disease Control and Prevention. *Enterobacter sakazakii* infections associated with the use of powdered infant formula—Tennessee, 2001. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5114a1.htm>. Accessed May 28, 2010.



Ingredients

Liquid Unflavored:

Water, and Casein Hydrolysate.

Allergens: Contains milk ingredients.

Availability

List Number	Item
62317	Liquid Protein Fortifier / 1.8-fl-oz (54-mL) Bottle / 48 ct

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Liquid Protein Fortifier

Extensively Hydrolyzed Protein

Nutrition Information - Liquid Unflavored

	6 mL
	Value
Calories	4
Protein, g	1
Water, g	5
Volume, mL	6

Preparation

Directions for Use

- Use as directed by health care professional.
- Shake gently.
- Mix with human milk or formula.
- Refrigerate fortifier after opening; use **within 24 hours or discard**,

Storage & Handling

Store unopened at room temperature; avoid extreme temperatures.

Product Information: Similac Expert Care® 24 Cal With Iron

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Expert Care® 24 Cal With Iron

Infant Formula with Iron

- When an iron-fortified feeding of increased caloric density is desired.
- Use under medical supervision.

Safety Precautions

- Never use a microwave oven to warm formula. Serious burns can result.



Ingredients

Liquid Unflavored:

Water, Nonfat Milk, Lactose, Soy Oil, Coconut Oil. Less than 0.5% of the Following: Ascorbic Acid, Monoglycerides, Soy Lecithin, Carrageenan, Choline Chloride, Ferrous Sulfate, Taurine, m-Inositol, dl-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Riboflavin, Calcium Pantothenate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Cyanocobalamin, Potassium Citrate, and Potassium Hydroxide.

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
51016	Similac Expert Care 24 Cal With Iron Institutional / 2-fl-oz (59-mL) Bottle / 48 ct

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Similac Expert Care® 24 Cal With Iron

Infant Formula with Iron

Nutrition Information - Liquid Unflavored

	100 Cal
	Value
Calories	100
Volume, mL	124
Protein, g	2.71
Fat, g	5.27
Carbohydrate, g	10.5
Water, g	109
Linoleic Acid, mg	1300
Vitamin A, IU	300
Vitamin D, IU	60
Vitamin E, IU	1.5
Vitamin K, mcg	8
Thiamin (Vitamin B1), mcg	100
Riboflavin (Vitamin B2), mcg	150
Vitamin B6, mcg	60
Vitamin B12, mcg	0.25
Niacin, mcg	1050
Folic Acid (Folacin), mcg	15
Pantothenic Acid, mcg	450
Biotin, mcg	4.4
Vitamin C (Ascorbic Acid), mg	9
Choline, mg	16
Inositol, mg	4.7
Calcium, mg	90
Phosphorus, mg	70
Magnesium, mg	7
Iron, mg	1.8
Zinc, mg	0.75
Manganese, mcg	5
Copper, mcg	90
Iodine, mcg	9
Selenium, mcg	1.8
Sodium, mg	32
Potassium, mg	132
Chloride, mg	81

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Expert Care® 24 Cal With Iron

Infant Formula with Iron

Preparation

Directions for Preparation and Use

- Do not add water.
- Shake very well.
- Do not use if breakaway ring is missing or broken.
- Ready to feed formula if mixed with water may not provide proper nutrition, and repeated use of such feedings could cause illness.
- Twist off cap.
- Twist on nipple and ring.
 - Use clean hands to attach nipple and ring.
- Once feeding begins, **use within one hour or discard.**

Storage & Handling

Ready to Feed

- Avoid prolonged exposure of bottles to light.
- Store unopened bottles at room temperature; avoid extreme temperatures.
- Do not reuse bottle.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Product Information: Similac Expert Care® Alimentum®

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Expert Care® Alimentum®

Hypoallergenic Infant Formula with Iron

- A nutritionally complete, hypoallergenic formula for infants, including those with colic symptoms due to protein sensitivity. A supplemental beverage for children with severe food allergies, sensitivity to intact protein, protein maldigestion, or fat malabsorption.
- Hypoallergenic; contains a predigested protein to virtually eliminate allergic reactions in most babies who are allergic to cow's milk protein.
- Hydrolyzed casein supplemented with free amino acids for infants who are sensitive to or unable to digest intact protein.
- Starts reducing colic symptoms due to protein sensitivity within 24 hours in most infants.^{*,1}
- DHA and ARA to help support brain and eye development.
- A unique blend of carbohydrates with two different absorptive pathways to help maximize absorption and minimize risk of malabsorption.
- Made with oils that have been shown to be well absorbed. Approximately 33% of the fat as medium chain triglycerides.
- Ready To Feed formula is corn-free.



Safety Precautions

- Never use a microwave oven to warm mixture. Serious burns can result.
- **Warning:** Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

^{*} Based on a clinical study with Similac Expert Care Alimentum Ready to Feed without DHA and ARA in a small group of infants experiencing colic symptoms due to cow's milk protein sensitivity.

¹ Data on File: Study AC84. August 2004. Abbott Nutrition, Columbus, Ohio.

Ingredients

Liquid Unflavored:

Water (87%), Sugar (4%), Casein Hydrolysate [Derived from Milk] (2%), Modified Tapioca Starch (2%), Safflower Oil (1.4%), Medium-Chain Triglycerides (1.3%), Soy Oil (1.1%). Less than 0.5% of the Following: C. Cohnii Oil, M. Alpina Oil, Calcium Citrate, Calcium Phosphate, Carrageenan, Potassium Phosphate, Ascorbic Acid, Magnesium Chloride, Calcium Hydroxide, Potassium Citrate, Salt, L-Cystine Dihydrochloride, Potassium Chloride, L-Tyrosine, Choline Chloride, L-Tryptophan, Ferrous Sulfate, Taurine, m-Inositol, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, L-Carnitine, Niacinamide, Calcium Pantothenate, Cupric Sulfate, Riboflavin, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Potassium Hydroxide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, and Cyanocobalamin.

Availability

List Number	Item
57508	Similac Expert Care Alimentum Ready to Feed / 8-fl-oz (237-mL) Can / Case of 24
57512	Similac Expert Care Alimentum Ready to Feed / 1-QT (946-mL) Bottle / Case of 6
57663	Similac Expert Care Alimentum Powder / 1-LB (454-g) SimplePac / Case of 6
59738	Similac Expert Care Alimentum Ready to Feed Institutional / 2-fl-oz (59-mL) Bottle / Case of 48

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Expert Care® Alimentum®

Hypoallergenic Infant Formula with Iron

Nutrition Information - Liquid Unflavored

	100 Cal	1000 mL
	Value	Value
Protein Equivalent, % Cal	11	11
Protein Equivalent Source	Casein Hydrolysate, L-Cystine, L-Tyrosine, L-Tryptophan	Casein Hydrolysate, L-Cystine, L-Tyrosine, L-Tryptophan
Fat, % Cal	48	48
Fat Source	Safflower Oil, Medium-Chain Triglycerides and Soy Oil (0.15% DHA, 0.40% ARA)	Safflower Oil, Medium-Chain Triglycerides and Soy Oil (0.15% DHA, 0.40% ARA)
Fat Ratio	38:33:28	38:33:28
Carbohydrate Source	Sugar, Modified Tapioca Starch	Sugar, Modified Tapioca Starch
Carbohydrate, % Cal	41	41
Carbohydrate Ratio	70:30	70:30
Osmolality, mOsm/kg H2O	370	370
Calories	100	676
Volume, mL	148	1000
Protein Equivalent, g	2.75	18.6
Fat, g	5.54	37.47
Carbohydrate, g	10.2	69.0
Water, g	133	899
Linoleic Acid, mg	1900	12,850
Potential Renal Solute Load, mOsm	25.3	171.3
Vitamin A, IU	300	2029
Vitamin D, IU	45	304
Vitamin E, IU	3.0	20.3
Vitamin K, mcg	15	101
Thiamin (Vitamin B1), mcg	60	406
Riboflavin, mcg	90	609
Vitamin B6, mcg	60	406
Vitamin B12, mcg	0.45	3.04
Niacin, mcg	1350	9130
Folic Acid (Folacin), mcg	15	101
Pantothenic Acid, mcg	750	5072
Biotin, mcg	4.5	30.4
Vitamin C (Ascorbic Acid), mg	9.0	61
Choline, mg	12	81
Inositol, mg	5	33.8
Calcium, mg	105	710
Calcium, mEq	5.2	35.4
Phosphorus, mg	75	507
Magnesium, mg	7.5	50.7
Iron, mg	1.8	12.17
Zinc, mg	0.75	5.07

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Similac Expert Care® Alimentum®

Hypoallergenic Infant Formula with Iron

Nutrition Information - Liquid Unflavored

	100 Cal	1000 mL
	Value	Value
Manganese, mcg	8	54
Copper, mcg	75	507
Iodine, mcg	15	101
Selenium, mcg	1.8	12.2
Sodium, mg	44	298
Sodium, mEq	1.9	12.9
Potassium, mg	118	798
Potassium, mEq	3.0	20.3
Chloride, mg	80	541
Chloride, mEq	2.3	15.5

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Similac Expert Care® Alimentum®

Hypoallergenic Infant Formula with Iron

Preparation

Ready to Feed

Your baby's health depends on carefully following these directions. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to boil (sterilize) bottles, nipples and rings before use.

2-fl-oz bottle

- Do not add water.
- Shake very well.
- **Do not use if breakaway ring is missing or broken.**
- Twist off cap.
- Twist on nipple and ring. Use clean hands to attach nipple and ring.
- Once feeding begins, **use within 1 hour or discard.**

8-fl-oz can

- Do not add water.
- Rinse can lid; shake very well.
- Lift and pull tab to open.
- Pour formula into bottle or cup; attach nipple or lid.
- Once feeding begins, **use within 1 hour or discard.**

1-QT bottle

- Do not add water.
- Shake well before each use.
- **Do not use if band around cap or inner foil seal is damaged.**
- Remove protective band; twist off and clean cap.
- Invert cap; press down to pierce foil, then turn cap a half turn.
- Remove foil.
- Pour formula into bottle; attach nipple.
- Once feeding begins, **use within 1 hour or discard.**
- Prepare and feed all formula in this container **within 2 days after opening.**

Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- Wash your hands, surfaces and utensils.
- Pour water into clean bottle (see mixing guide).
- Add 1 unpacked level scoop (8.7 g) to each 2 fl oz of water.
- Return dry scoop to can.
- Cap bottle; shake well; attach nipple.
- Once feeding begins, **use within 1 hour or discard.**

Powder Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.7 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz
8 fl oz		4 scoops		8 fl oz

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Storage & Handling

Ready to Feed

2-fl-oz bottle

- Avoid prolonged exposure to light.
- Store unopened bottles at room temperature; avoid extreme temperatures.

8-fl-oz can

- Cover opened cans and prepared bottles not used immediately. Store in refrigerator and **use within 48 hours.**
- Store unopened bottles at room temperature; avoid extreme temperatures.

1-QT bottle

- Once opened, store quart bottle in refrigerator.
- Store prepared bottles in refrigerator and **feed to baby within 48 hours.**
- Store unopened containers at room temperature; avoid extreme temperatures.
- Do not reuse container.
- Prepare and feed all formula from this container **within 2 days after opening.**

Powder

- Once mixed, store bottles in refrigerator and **feed to baby within 24 hours.**
- Store unopened or opened container at room temperature; avoid extreme temperatures.
- **Use opened can contents within 1 month.**

Product Information: Similac Expert Care® for Diarrhea

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Expert Care® for Diarrhea

Infant formula for the dietary management of diarrhea

- For dietary management of diarrhea to help firm loose and watery stools in infants older than 6 months and toddlers.
- First and only nutritionally complete infant formula to contain added dietary fiber (soy) specifically for dietary management of diarrhea (6 g/L).
- Clinically shown to reduce the duration of loose, watery stools during mild to severe diarrhea in older infants.
- Clinically shown to be effective in the management of antibiotic-induced diarrhea.
- A unique blend of two carbohydrates using two absorptive pathways to help maximize absorption and minimize malabsorption risks.
- Low osmolality (240 mOsm/kg water) to reduce the risk of osmotic diarrhea.
- Lactose-free.
- Gluten-free.
- Kosher.
- Halal.



Safety Precautions

- **Never use a microwave oven to warm formula.** Serious burns can result.
- **This product should not be fed to infants and toddlers with constipation.**

Ingredients

Liquid Unflavored:

Water (86%), Corn Syrup (5%), Sugar (3%), Soy Oil (2%), Soy Protein Isolate (2%), Coconut Oil (1.4%), Soy Fiber (0.77%). Less than 0.5% of the Following: Calcium Citrate, Potassium Citrate, Calcium Phosphate, Potassium Phosphate, Potassium Chloride, Monoglycerides, Soy Lecithin, Magnesium Chloride, Carrageenan, Salt, Ascorbic Acid, Choline Chloride, L-Methionine, Taurine, Ferrous Sulfate, m-Inositol, Zinc Sulfate, d-Alpha-Tocopheryl Acetate, L-Carnitine, Niacinamide, Calcium Pantothenate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Potassium Hydroxide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, and Cyanocobalamin.

Allergens: Contains soy ingredients.

Availability

List Number	Item
51276	Similac Expert Care Diarrhea Ready to Feed / 8-fl-oz (237-mL) can / 24 ct
57768	Similac Expert Care Diarrhea Ready to Feed / 1-QT (946-mL) Bottle / 6 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Expert Care® for Diarrhea

Infant formula for the dietary management of diarrhea

Nutrition Information - Liquid Unflavored

	100 Cal	1000 mL
	Value	Value
Protein, % Cal	11	11
Protein Source	Soy Protein Isolate, L-Methionine	Soy Protein Isolate, L-Methionine
Fat, % Cal	49	49
Fat Source	Soy and Coconut Oils	Soy and Coconut Oils
Fat Ratio	60:40	60:40
Carbohydrate, % Cal	40	40
Carbohydrate Source	Corn Syrup Solids, Sugar	Corn Syrup Solids, Sugar
Carbohydrate Ratio	60:40	60:40
Osmolality, mOsm/kg H2O	240	240
Calories	100	676
Volume, mL	148	1000
Protein, g	2.66	17.99
Fat, g	5.46	36.93
Carbohydrate, g	10.1	68.3
Water, g	133	899
Linoleic Acid, mg	1300	8792
Potential Renal Solute Load, mOsm	24.0	162.6
Vitamin A, IU	300	2029
Vitamin D, IU	60	406
Vitamin E, IU	1.5	10.1
Vitamin K, mcg	11	74
Thiamin (Vitamin B1), mcg	60	406
Riboflavin (Vitamin B2), mcg	90	609
Vitamin B6, mcg	60	406
Vitamin B12, mcg	0.45	3.04
Niacin, mcg	1350	9130
Folic Acid, mcg	15	101
Pantothenic Acid, mcg	750	5072
Biotin, mcg	4.5	30.4
Vitamin C, mg	9	61
Choline, mg	12	81
Inositol, mg	5	33.8
Calcium, mg	105	710
Calcium, mEq	5.2	35.4
Phosphorus, mg	75	507
Magnesium, mg	7.5	50.7
Iron, mg	1.8	12.2
Zinc, mg	0.75	5.07
Manganese, mcg	25	169

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Similac Expert Care® for Diarrhea

Infant formula for the dietary management of diarrhea

Nutrition Information - Liquid Unflavored

	100 Cal	1000 mL
	Value	Value
Copper, mcg	75	507
Iodine, mcg	15	101
Selenium, mcg	1.8	12.2
Sodium, mg	44	298
Sodium, mEq	1.9	12.9
Potassium, mg	108	730
Potassium, mEq	2.8	18.7
Chloride, mg	62	419
Chloride, mEq	1.8	11.8

Preparation

Ready-to-Feed

Your baby's health depends on carefully following these directions. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to boil (sterilize) bottles, nipples and rings before use.

8-fl-oz can

- **Do not add water.**
- Rinse can lid; shake very well.
- Lift and pull tab to open.
- Pour formula into bottle or cup; attach nipple or lid; once feeding begins, **use within 1 hour or discard.**

1-QT bottle

- **Do not add water.**
- **Do not use if band around cap or inner foil seal is damaged.**
- Shake very well before each use.
- Remove protective band; twist off and clean cap.
- Invert cap; press down to pierce foil, then turn cap a half turn.
- Remove foil.
- Pour formula into bottle or cup; attach nipple or lid.
- Once feeding begins, **use within 1 hour or discard.**

Storage & Handling

Ready-to-Feed

8-fl-oz can

- Cover opened cans and prepared bottles not used immediately.
- Store in refrigerator and **use within 48 hours.**
- Store unopened cans at room temperature; avoid extreme temperatures.

1-QT bottle

- Once opened, store quart bottle in refrigerator.
- Store prepared bottles in refrigerator and **feed to baby within 48 hours.**
- Store unopened containers at room temperature; avoid extreme temperatures.
- Prepare and feed all formula from this container **within 3 days after opening.**
- Do not reuse container.

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Product Information: Similac Expert Care® NeoSure®

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Expert Care® NeoSure®

Infant Formula with Iron

- Similac Expert Care NeoSure is designed for babies who were born prematurely.
- Use under medical supervision.
- Studies have shown that preterm babies fed Similac Expert Care NeoSure had: excellent catch-up growth^{*,†}; improved language development^{†,‡,2}; and improved visual development.^{†,§,2}
- 25% of the fat blend as medium-chain triglycerides, an easily digested and well-absorbed fat source.
- Has a unique blend of nucleotides to help support the developing immune system.^{3,4,5,6}
- Lutein and DHA to support the developing eyes and brain.
- Halal.
- Kosher.



Safety Precautions

- Never use a microwave to warm formula. Serious burns can result.
- **Warning:** Powdered infant formulas are not sterile and may be fed to premature infants or infants who have immune problems only as directed by your baby's doctor.

^{*} When compared to premature infants fed term infant formula.

[†] Compared to infants fed a formula without DHA and ARA in a clinical trial with Similac Special Care and Similac Expert Care NeoSure Infant Formula with Iron.

[‡] Based on a post-hoc analysis of English-speaking singleton premature infants using MacArthur Communicative Developmental Inventories.

[§] Visual acuity measured at 4 & 6 months corrected age and assessed by Visual Evoked Potential (VEP).

¹ Carver JD, et al. *Pediatrics* 2001;107:683-689.

² O'Connor DL, Hall R, Adamkin D, et al. *Pediatrics* 2001;108:359-371.

³ Buck, RH, et al. *Pediatr Res* 2004; 56:891-900.

⁴ Schaller, JP, et al. *Pediatr Res* 2004; 56:883-890.

⁵ Pickering KL, et al. *Pediatrics* 1998; 101:242-249.

⁶ Yau K, et al. *J Pediatr Gastroenterol Nutr* 2003; 36:37-43.

Ingredients

Powder Unflavored:

Nonfat Milk, Corn Syrup Solids, Lactose, Soy Oil, High Oleic Safflower Oil, Whey Protein Concentrate, Medium-Chain Triglycerides, Coconut Oil. Less than 2% of the Following: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Potassium Citrate, Calcium Phosphate, m-Inositol, Ascorbic Acid, Magnesium Chloride, Soy Lecithin, Calcium Carbonate, Taurine, Ferrous Sulfate, Choline Bitartrate, Choline Chloride, Ascorbyl Palmitate, L-Carnitine, Potassium Chloride, Salt, Zinc Sulfate, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Sodium Citrate, Niacinamide, Potassium Phosphate, Calcium Pantothenate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Cyanocobalamin, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
56177	Similac Expert Care NeoSure Ready to Feed Institutional / 2-fl-oz (59-mL) Bottle / 48 ct
57430	Similac Expert Care NeoSure Powder / 13.1-oz (371-g) Can / 6 ct
57455	Similac Expert Care NeoSure Ready to Feed / 1-QT (946-mL) Bottle / 6 ct
59645	Similac Expert Care NeoSure Ready to Feed / 2-fl-oz (59-mL) Bottle/ 6 x 8 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Powder Unflavored

	100 Cal*	1000 mL*
	Value	Value
Calories	100	744
Volume, mL	134	1000
Protein, g	2.8	20.83
Fat, g	5.5	40.92
Carbohydrate, g	10.1	75.1
Water, g	120	893
Linoleic Acid, mg	750	5579
Vitamin A, IU	350	2604
Vitamin D, IU	70	521
Vitamin E, IU	3.6	26.8
Vitamin K, mcg	11	81.8
Thiamin (Vitamin B1), mcg	175	1302
Riboflavin (Vitamin B2), mcg	150	1116
Vitamin B6, mcg	100	744
Vitamin B12, mcg	0.40	2.98
Niacin, mcg	1950	14506
Folic Acid (Folacin), mcg	25	186
Pantothenic Acid, mcg	800	5951
Biotin, mcg	9	67.0
Vitamin C (Ascorbic Acid), mg	15	112
Choline, mg	16	119
Inositol, mg	35	260
Calcium, mg	105	781
Phosphorus, mg	62	461
Magnesium, mg	9	67.0
Iron, mg	1.8	13.4
Zinc, mg	1.2	8.9
Manganese, mcg	10	74
Copper, mcg	120	893
Iodine, mcg	15	112
Selenium, mcg	2.3	17.1
Sodium, mg	33	245
Potassium, mg	142	1056
Chloride, mg	75	558

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Powder Unflavored Footnotes & References

Per 100 Cal*

*Nutrient values are applicable when prepared as directed.

Per 1000 mL*

*Nutrient values are applicable when prepared as directed.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Preparation

Ready to Feed

Your baby's health depends on carefully following these directions. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to boil (sterilize) bottles, nipples and rings before use. It is important to consult your baby's doctor before changing formula.

2-fl-oz bottle

- **Do not use if breakaway ring is missing or broken.**
- Do not add water. Ready to feed formula if mixed with water may not provide proper nutrition and repeated use of such feedings could cause illness.
- Shake very well.
- Twist off cap.
- Twist on nipple and ring.
 - Since this formula is commercially sterile, use clean hands to attach any clean standard nipple and ring.
- No need to warm formula before feeding.
- Once feeding begins, **use within 1 hour or discard.**

1-QT bottle

- **Do not use if band around cap or inner foil seal is damaged.**
- Do not add water.
- Shake very well before each use.
- Remove protective band, twist off and clean cap.
- Invert cap; press down to pierce foil, then turn cap a half turn.
- Remove foil.
- Pour formula into bottle; attach nipple.
- Once feeding begins, **use within 1 hour or discard.**
- Prepare and feed all formula from the 1 QT container **within 3 days after opening.**

Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottle, nipple and ring before use.

- Wash your hands, surfaces and utensils.
- Pour desired amount of water into clean bottle (see mixing guide).
- Add 1 unpacked level scoop (9.6 g) to each 2 fl oz of water.
- Return dry scoop to can.
- Cap bottle; shake well; attach nipple.
- Once feeding begins, **use within 1 hour or discard.**

Powder Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (9.6 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
8 fl oz		4 scoops		8 fl oz

Storage & Handling

Ready to Feed

2-fl-oz bottle

- Store unopened at room temperature; avoid extreme temperatures.
- Do not reuse this bottle.
- **Protect from light.**
- **Store in carton until ready for use.**

1-QT bottle

- Once opened, store quart bottle in refrigerator.
- Once prepared, store bottles in refrigerator and **feed to baby within 48 hours.**
- Store unopened containers at room temperature; avoid extreme temperatures.
- **Use within 3 days after opening.**
- Do not reuse container.

Powder

- Once mixed, store bottles in refrigerator and **feed to baby within 24 hours.**
- Store unopened or opened container at room temperature; avoid extreme temperatures.
- **Use opened can contents within 1 month.**

Product Information: Similac For Spit-Up®

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac For Spit-Up®

Infant Formula with Iron

- A nutritionally complete milk-based formula with added rice starch to help reduce frequent spit up.
- Milk-based, reduced-lactose formula* suitable for lactose sensitivity.
- Our exclusive formula has:
 - Lutein and DHA for baby's developing eyes and brain.
 - Calcium for strong bones - no palm olein oil.
 - Nucleotides to help support the immune system.
 - Prebiotics to help promote digestive health.
 - Carotenoids naturally found in breast milk.
- Easy to digest milk-based, reduced lactose formula with added rice starch.
- Clinically shown to reduce frequent spit up in healthy babies by 54%.^{1,†}
- Clinically shown to support normal infant growth.
- Reduced lactose formulation suitable for infants with lactose sensitivity.
- Kosher.
- Halal.
- Gluten-free.



Safety Precautions

- **Never use a microwave oven to warm formula.** Serious burns can result.
- Not for use by infants or children with galactosemia.
- **Warning:** Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

* Compared to standard milk-based formulas.

† Among healthy 2-month-old infants compared to a standard formula.

¹ Data on file, Study AJ68, May 2007. Abbott Nutrition, Columbus, Ohio.

Ingredients

Powder Unflavored:

Corn Syrup Solids, Rice Starch, Milk Protein Isolate, High Oleic Safflower Oil, Sugar, Soy Oil, Coconut Oil, Galactooligosaccharides. Less than 2% of the Following: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Lycopene, Calcium Phosphate, Potassium Citrate, Potassium Chloride, Potassium Phosphate, Calcium Carbonate, Magnesium Chloride, Ascorbic Acid, L-Cystine Dihydrochloride, Choline Chloride, Ascorbyl Palmitate, Salt, Ferrous Sulfate, Choline Bitartrate, Taurine, m-Inositol, dl-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Zinc Sulfate, L-Carnitine, Niacinamide, Calcium Pantothenate, Vitamin A Palmitate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Cyanocobalamin, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).

Allergens: Contains milk ingredients.

Availability

List Number	Item
50959	Similac For Spit-Up Powder / 12.3-oz (349-g) Can / 6 ct
53729	Similac for Spit Up Powder / 1.41-LB (638-g) SimplePac / 6 ct
56728	Similac for Spit Up Ready to Feed / 1-QT (946-mL) Bottle / 6 ct
62139	Similac for Spit Up Ready to Feed / 2-fl-oz (59-mL) Bottle / 48 ct

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Nutrition Information - Powder Unflavored

	100 Cal*	1000 mL*
	Value	Value
Calories	100	676
Volume, mL	148	1000
Protein, g	2.14	14.47
Fat, g	5.64	38.1
Carbohydrate, g	10.40	70.3
Water, g	133	899
Linoleic Acid, mg	1000	6760
Potential Renal Solute Load, mOsm	19.9	134.7
Vitamin A, IU	300	2029
Vitamin D, IU	60	406
Vitamin E, IU	3.0	20.3
Vitamin K, mcg	8	54
Thiamin (Vitamin B1), mcg	100	676
Riboflavin (Vitamin B2), mcg	150	1014
Vitamin B6, mcg	60	406
Vitamin B12, mcg	0.25	1.7
Niacin, mcg	1050	7101
Folic Acid (Folacin), mcg	15	101
Pantothenic Acid, mcg	450	3043
Biotin, mcg	4.4	29.8
Vitamin C (Ascorbic Acid), mg	9	61
Choline, mg	16	108
Inositol, mg	4.3	29.1
Calcium, mg	84	568
Calcium, mEq	4.2	28.3
Phosphorus, mg	56	379
Magnesium, mg	6	40.6
Iron, mg	1.8	12.2
Zinc, mg	0.75	5.1
Manganese, mcg	5	34
Copper, mcg	90	609
Iodine, mcg	9	61
Selenium, mcg	1.8	12.2
Sodium, mg	30	203
Sodium, mEq	1.3	8.8
Potassium, mg	107	724
Potassium, mEq	2.7	18.5
Chloride, mg	65	440
Chloride, mEq	1.8	12.4

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Powder Unflavored Footnotes & References

Per 100 Cal*

*Nutrient values are applicable when prepared as directed.

Per 1000 mL*

*Nutrient values are applicable when prepared as directed.

Preparation

Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- Wash your hands, surfaces and utensils.
- Pour water into clean bottle (see mixing guide).
- Add 1 unpacked level scoop (8.7 g) to each 2 fl oz of water.
- Return dry scoop to container.
- Cap bottle; shake well; wait two minutes; attach nipple; gently swirl before feeding.
- Once feeding begins, use **within 1 hour or discard**.

Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.7 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz
8 fl oz		4 scoops		8 fl oz

Ready to Feed

Your baby's health depends on carefully following these directions. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to boil (sterilize) bottles, nipples and rings before use.

2-fl-oz Bottle

- Do not add water.
- Shake very well.
- **Do not use if breakaway ring is missing or broken.**
- Twist off cap.
- Twist on nipple and ring; use clean hands to attach nipple and ring.
- Once feeding begins, use **within 1 hour or discard**.

1-Qt Bottle

- Do not add water.
- Shake very well before each use.
- Remove protective band; twist off and clean cap. **Do not use if band around cap or inner foil seal is damaged.**
- Invert cap; press down to piece foil, then turn cap a half turn.
- Remove foil.
- Pour formula into bottle; attach nipple.
- Once feeding begins, use **within 1 hour or discard**.
- Prepare and feed all formula from the container **within 3 days after opening**.

Storage & Handling

Powder

- Once mixed, store bottles in refrigerator and **feed to baby within 24 hours**.
- Store unopened or opened container at room temperature; avoid extreme temperatures.
- Use opened can contents **within 1 month**.
- Do not reuse container.

Ready to Feed

2-fl-oz Bottle

- Avoid prolonged exposure of bottles to light.
- Store unopened containers at room temperature; avoid extreme temperatures.

1-QT Bottle

- Once opened, store quart bottle in refrigerator.
- Store prepared bottles in refrigerator and **feed to baby within 48 hours**.
- Store unopened containers at room temperature; avoid extreme temperatures.
- Prepare and feed all formula from this container **within 3 days after opening**.
- Do not reuse container.

Product Information: Similac Go & Grow® Milk-Based Formula

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Go & Grow® Milk-Based Formula

Milk-Based Infant Formula with Iron

- A milk-based, iron-fortified formula for older babies 9 to 24 months old. Specially formulated to help bridge the nutritional gaps that can be associated with the transition to table foods.
- Our exclusive formula has:
 - DHA (Omega 3) and ARA (Omega 6) for brain and eye development.
 - Calcium for strong bones - no palm olein oil.
 - Nucleotides to help support the immune system.
 - Prebiotics to help promote digestive health.
 - Carotenoids naturally found in breast milk.
 - Lutein to support eye health.
- Over 25 essential minerals and vitamins.
- At least 30% of the recommended Daily Value* of calcium, iron, vitamin C and vitamin E.
- Complements nutrition provided by table food.
- Gluten-free.
- Kosher.
- Halal.



Safety Precautions

- Never use a microwave oven to warm formula. Serious burns can result.
- **Warning:** Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

* Percent daily value (DV) for 1- to 4-year-olds

Ingredients

Powder Unflavored:

Nonfat Milk, Lactose, High Oleic Safflower Oil, Soy Oil, Coconut Oil, Galactooligosaccharides. Less than 2% of the Following: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Lycopene, Calcium Phosphate, Potassium Citrate, Ascorbic Acid, Soy Lecithin, Calcium Carbonate, Choline Chloride, Ferrous Sulfate, Choline Bitartrate, Ascorbyl Palmitate, Taurine, m-Inositol, d-Alpha-Tocopheryl Acetate, Zinc Sulfate, Mixed Tocopherols, Niacinamide, Calcium Pantothenate, Vitamin A Palmitate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Cyanocobalamin, Potassium Phosphate, Magnesium Chloride, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
50827	Similac Go & Grow Milk-Based EarlyShield Formula Powder / 1.38-lb (624-g) SimplePac / 6 ct

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Similac Go & Grow® Milk-Based Formula

Milk-Based Infant Formula with Iron

Nutrition Information - Powder Unflavored

	100 Cal [*]	1000 mL [*]
	Value	Value
Calories	100	650
Volume, mL	154	1000
Protein, g	3.00	19.51
Fat, g	5.40	35.12
Carbohydrate, g	10.2	66.3
Water, g	133	899
Linoleic Acid, mg	1000	6503
Vitamin A, IU	300	1951
Vitamin D, IU	60	390
Vitamin E, IU	3	19.5
Vitamin K, mcg	8	52
Thiamin (Vitamin B1), mcg	100	650
Riboflavin (Vitamin B2), mcg	150	975
Vitamin B6, mcg	60	390
Vitamin B12, mcg	0.3	1.95
Niacin, mcg	1050	6828
Folic Acid, mcg	15	98
Pantothenic Acid, mcg	450	2926
Biotin, mcg	4.4	28.6
Vitamin C, mg	12	78
Choline, mg	16	104
Inositol, mg	4.7	31
Calcium, mg	195	1268
Phosphorus, mg	130	845
Magnesium, mg	9	58.5
Iron, mg	2	13.0
Zinc, mg	0.75	4.88
Manganese, mcg	5	33
Copper, mcg	90	585
Iodine, mcg	12	78
Selenium, mcg	1.8	11.7
Sodium, mg	30	195
Potassium, mg	150	975
Chloride, mg	80	520

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Go & Grow® Milk-Based Formula

Milk-Based Infant Formula with Iron

Powder Unflavored Footnotes & References

Per 100 Cal*

*Nutrient values are applicable when prepared as directed.

Per 1000 mL*

*Nutrient values are applicable when prepared as directed.

Preparation

Powder: Caloric Density

The following table shows the quantity of water to mix with the number of **unpacked, level scoop(s) of Powder** to arrive at the approximate caloric densities shown. Use only the scoop provided in the container.

Caloric Density (Cal/fl oz)	Water (fl oz)	Unpacked, Level Scoop	Approximate Yield (fl oz)
20 (standard)	2	1	2
22	3.5	2	4
24	5	3	6
26	1.5	1	2
27	4.25	3	5

Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- Wash your hands, surfaces and utensils.
- Pour water into a clean cup or bottle (see mixing guide).
- Add 1 unpacked level scoop (8.7 g) to each 2 fl oz of water.
- Return dry scoop to holder in lid.
- Cap cup or bottle.
- Shake well.
- Once feeding begins, **use within 1 hour or discard.**

Standard Preparation Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.7 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz
8 fl oz		4 scoops		8 fl oz

Pour prepared formula into feeding cup or bottle, cap and store in refrigerator. **Use within 24 hours.**

Similac Go & Grow® Milk-Based Formula

Milk-Based Infant Formula with Iron

Storage & Handling

Powder

- Once mixed, store bottle or cup in refrigerator and **feed within 24 hours**.
- Store unopened or opened container at room temperature; avoid extreme temperatures.
- **Use opened container contents within 1 month.**
- Do not reuse container.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Product Information: Similac Go & Grow® Soy-Based Formula

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Go & Grow® Soy-Based Formula

Soy-Based Infant Formula with Iron

- A soy-based, iron-fortified formula for older babies and toddlers 9 to 24 months old. Specially formulated to help bridge the nutritional gaps that can be associated with the transition to table foods. A lactose-free feeding for older babies when lactose should be avoided, including lactase deficiency, lactose intolerance, and galactosemia.
- Milk-free, lactose-free nutritional formula.
- Complements nutrition provided by table food.
- Soy protein isolate to help manage IgE-mediated cow's-milk-protein allergy or sensitivity.
- A unique blend of two carbohydrates—corn syrup and sucrose—using two absorptive pathways to help maximize absorption and minimize malabsorption risks.
- Lutein and DHA to support the developing eyes and brain.
- Over 25 essential minerals and vitamins in 8 fl oz.
- 30% Daily Value* for calcium in 8 fl oz for strong bones.
- 32% Daily Value* for iron in 8 fl oz for brain development.
- 48% Daily Value* for vitamin C in 8 fl oz, an important antioxidant.
- Halal.
- Kosher.



Safety Precautions

- Never use a microwave to warm formula. Serious burns can result.
- **Warning:** Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

* Percent daily value (DV) for 1- to 4-year-olds

Ingredients

Powder Unflavored:

Corn Syrup (38%), Soy Protein Isolate (16%), High Oleic Safflower Oil (11%), Sugar (10%), Soy Oil (8%), Coconut Oil (7%), Calcium Phosphate (3%). Less than 2% of the Following: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Lycopene, Fructooligosaccharides, Potassium Citrate, Salt, Magnesium Chloride, Ascorbic Acid, L-Methionine, Potassium Chloride, Choline Chloride, Taurine, Ferrous Sulfate, Ascorbyl Palmitate, m-Inositol, Mixed Tocopherols, Zinc Sulfate, d-Alpha-Tocopheryl Acetate, L-Carnitine, Niacinamide, Calcium Carbonate, Calcium Pantothenate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Phylloquinone, Biotin, Vitamin D3, Cyanocobalamin, Sodium Selenate, and Potassium Hydroxide.

Allergens: Contains soy ingredients.

Availability

List Number	Item
50837	Similac Go & Grow Soy-Based EarlyShield Powder / 1.38-LB (624-g) SimplePac / 6 ct

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Similac Go & Grow® Soy-Based Formula

Soy-Based Infant Formula with Iron

Nutrition Information - Powder Unflavored

	100 Cal [*]	1000 mL [*]
	Value	Value
Calories	100	676
Volume, mL	148	1000
Protein, g	2.80	18.93
Fat, g	5.40	36.89
Carbohydrate, g	10.3	69.6
Water, g	133	899
Linoleic Acid, mg	1000	6757
Vitamin A, IU	300	2029
Vitamin D, IU	60	406
Vitamin E, IU	3	20.3
Vitamin K, mcg	11	74
Thiamin (Vitamin B1), mcg	60	406
Riboflavin (Vitamin B2), mcg	90	609
Vitamin B6, mcg	60	406
Vitamin B12, mcg	0.45	3.04
Niacin, mcg	1350	9130
Folic Acid, mcg	15	101
Pantothenic Acid, mcg	750	5072
Biotin, mcg	4.5	30.4
Vitamin C, mg	12	81
Choline, mg	12	81
Inositol, mg	5	33.8
Calcium, mg	195	1318
Phosphorus, mg	130	879
Magnesium, mg	7.5	50.7
Iron, mg	2.0	13.5
Zinc, mg	0.75	5.07
Manganese, mcg	25	169
Copper, mcg	75	507
Iodine, mcg	15	101
Selenium, mcg	1.8	12.2
Sodium, mg	44	298
Potassium, mg	120	811
Chloride, mg	62	419

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Go & Grow® Soy-Based Formula

Soy-Based Infant Formula with Iron

Powder Unflavored Footnotes & References

Per 100 Cal*

*Nutrient values are applicable when prepared as directed.

Per 1000 mL*

*Nutrient values are applicable when prepared as directed.

Preparation

Powder: Caloric Density

The following table shows the quantity of water to mix with the number of **unpacked, level scoop(s) of Powder** to arrive at the approximate caloric densities shown. Use only the scoop provided in the container.

Caloric Density (Cal/fl oz)	Water (fl oz)	Unpacked, Level Scoop	Approximate Yield (fl oz)
20 (standard)	2	1	2
22	3.5	2	4
24	5	3	6
26	1.5	1	2
27	4.25	3	5

To make a larger amount of formula at standard density (20 Cal/fl oz) using a standard measuring cup, add **1 unpacked, level cup (NOT the enclosed scoop) of Powder (100 g)** to 23 fl oz of water. Yields approximately 26 fl oz of formula. Use within 24 hours.

Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- Wash your hands, surfaces and utensils.
- Pour water into a clean cup or bottle (see mixing guide).
- Add 1 unpacked level scoop (8.7 g) to each 2 fl oz of water.
- Return dry scoop to holder in lid.
- Cap cup or bottle.
- Shake well.
- Once feeding begins, **use within 1 hour or discard.**

Standard Preparation Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.7 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz
8 fl oz		4 scoops		8 fl oz

Pour prepared formula into feeding cup or bottle, cap and store in refrigerator. **Use within 24 hours.**

Storage & Handling

Powder

- Once mixed, store bottles in refrigerator and **feed within 24 hours.**
- Store unopened or opened container at room temperature; avoid extreme temperatures.
- **Use opened container contents within 1 month.**
- Do not reuse container.

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Product Information: Similac Sensitive®

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Sensitive®

Infant Formula with Iron

- A nutritionally complete, reduced-lactose infant feeding* that is an alternative to standard milk-based formulas for mild tolerance symptoms such as fussiness and gas due to lactose sensitivity.
- Our exclusive formula has:
 - Lutein and DHA for baby's developing eyes and brain.
 - Calcium for strong bones - no palm olein oil.
 - Nucleotides to help support the immune system.
 - Prebiotics to help promote digestive health.
 - Carotenoids naturally found in breast milk.
- Clinically shown to support normal infant growth.
- A unique blend of two carbohydrates to help maximize absorption and minimize malabsorption risks.
- Reduced lactose formulation suitable for infants with lactose sensitivity.
- Low osmolality (200 mOsm/kg water).
- Kosher.
- Halal.
- Gluten-free.



Safety Precautions

- **Never use a microwave oven to warm formula.** Serious burns can result.
- Not for use by infants or children with galactosemia.
- **Warning:** Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.
- **Hypo- and hypercaloric formulas should be used under the direction of a physician.**
- 27 Cal/fl oz or more calorically dense formula may not supply enough water for some infants. Hydration status should be monitored and water supplied from other sources if necessary.

* Compared to standard milk-based formulas.

Ingredients

Powder Unflavored:

Corn Syrup, Sugar, Milk Protein Isolate, High Oleic Safflower Oil, Soy Oil, Coconut Oil, Galactooligosaccharides. Less than 2% of the Following: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Lycopene, Calcium Phosphate, Potassium Citrate, Potassium Chloride, Sodium Citrate, Magnesium Phosphate, Ascorbic Acid, Calcium Carbonate, Choline Chloride, Ferrous Sulfate, Magnesium Chloride, Ascorbyl Palmitate, Choline Bitartrate, Taurine, m-Inositol, dl-Alpha-Tocopheryl Acetate, Zinc Sulfate, Mixed Tocopherols, L-Carnitine, Niacinamide, Calcium Pantothenate, Vitamin A Palmitate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Cyanocobalamin, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).

Allergens: Contains milk ingredients.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Availability

List Number	Item
50817	Similac Sensitive Powder / 1.41-lb (638-g) SimplePac / 6 ct
53445	Similac Sensitive Powder / 2.06-lb (936-g) SimplePac / 6 ct
53676	Similac Sensitive On-the-go Ready to Feed / 8-fl-oz (237-mL) Bottle / 4 x 6 ct
53907	Similac Sensitive Ready to Feed 6-Pack / 1-QT (946-mL) Bottle
56170	Similac Sensitive Ready to Feed 8-Pack / 1-QT (946-mL) Bottle
56258	Similac Sensitive Concentrated Liquid / 1-QT (946-mL) Bottle / 6 ct
56508	Similac Sensitive Ready to Feed Institutional / 2-fl-oz (59-mL) Bottle / 48 ct
57533	Similac Sensitive Ready to Feed / 1-QT (946-mL) Bottle / 6 ct
57535	Similac Sensitive Concentrated Liquid / 13-fl-oz (384-mL) Can / 12 ct
57539	Similac Sensitive EarlyShield Powder / 12.6-oz (357-g) Can / 6 ct
62951	Similac Sensitive Powder / 1.86-lb (845-g) SimplePac / 6 ct

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Nutrition Information - Powder Unflavored

	100 Cal*	1000 mL*
	Value	Value
Protein, % Cal	9	9
Protein Source	Milk Protein Isolate	Milk Protein Isolate
Fat, % Cal	49	49
Fat Source	High Oleic Safflower, Soy and Coconut Oil (0.15% DHA, 0.40% ARA)	High Oleic Safflower, Soy and Coconut Oil (0.15% DHA, 0.40% ARA)
Fat Ratio	40:30:29	40:30:29
Carbohydrate, % Cal	43	43
Carbohydrate Source	Corn Maltodextrin, Sugar, Galactooligosaccharides (GOS)	Corn Maltodextrin, Sugar, Galactooligosaccharides (GOS)
Prebiotic†	GOS	GOS
Carbohydrate Ratio	53:43:4	53:43:4
Osmolality, mOsm/kg H2O	200	200
Calories	100	676
Volume, mL	148	1000
Protein, g	2.14	14.47
Fat, g	5.4	36.52
Carbohydrate, g	11.1	75.1
Water, g	133	899
Linoleic Acid, mg	1000	6763
Potential Renal Solute Load, mOsm	19.9	134.7
Vitamin A, IU	300	2029
Vitamin D, IU	60	406
Vitamin E, IU	3.0	20.3
Vitamin K, mcg	8	54
Thiamin (Vitamin B1), mcg	100	676
Riboflavin (Vitamin B2), mcg	150	1014
Vitamin B6, mcg	60	406
Vitamin B12, mcg	0.25	1.69
Niacin, mcg	1050	7101
Folic Acid (Folacin), mcg	15	101
Pantothenic Acid, mcg	450	3043
Biotin, mcg	4.4	29.8
Vitamin C (Ascorbic Acid), mg	9	61
Choline, mg	16	108
Inositol, mg	4.3	29.1
Calcium, mg	84	568
Calcium, mEq	4.2	28.3
Phosphorus, mg	56	379
Magnesium, mg	6	40.6

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Nutrition Information - Powder Unflavored

	100 Cal [*]	1000 mL [*]
	Value	Value
Iron, mg	1.8	12.2
Zinc, mg	0.75	5.07
Manganese, mcg	5	34
Copper, mcg	90	609
Iodine, mcg	9	61
Selenium, mcg	1.8	12.2
Sodium, mg	30	203
Sodium, mEq	1.3	8.8
Potassium, mg	107	724
Potassium, mEq	2.7	18.5
Chloride, mg	65	440
Chloride, mEq	1.8	12.4

Powder Unflavored Footnotes & References

Per 100 Cal^{*}

^{*}Nutrient values are applicable when prepared as directed.

[†]GOS is approximately 3% of the total carbohydrate and provides approximately 1.3 Cal/g.

Per 1000 mL^{*}

^{*}Nutrient values are applicable when prepared as directed.

[†]GOS is approximately 3% of the total carbohydrate and provides approximately 1.3 Cal/g.

Preparation

Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- Wash your hands, surfaces and utensils.
- Pour desired amount of water into clean bottle (see mixing guide).
- Add 1 unpacked level scoop (8.8 g) to each 2 fl oz of water.
- Return dry scoop to container.
- Cap bottle; shake well; attach nipple.
- Once feeding begins, **use within 1 hour or discard**.

Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.8 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

8 fl oz		4 scoops		8 fl oz
---------	--	----------	--	---------

Caloric Density

The following table shows the quantity of water to mix with the number of **unpacked, level scoop(s) of Powder** to arrive at the approximate caloric densities shown. Use only the scoop provided in the container.

Caloric Density (Cal/fl oz)	Water (fl oz)	Unpacked, Level Scoop	Approximate Yield (fl oz)
20 (standard)	2	1	2
22	3.5	2	4
24	5	3	6
26	1.5	1	2
27	4.25	3	5

To make a larger amount of formula at standard density (20 Cal/fl oz) using a standard measuring cup, add **1 unpacked level cup (NOT the enclosed scoop) of Powder (100 g)** to 23 fl oz of water. Yields approximately 26 fl oz of formula. Prepared formula should not be left unrefrigerated. Use within 24 hours.

Concentrated Liquid

Your baby's health depends on carefully following these directions. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and need to boil (sterilize) bottles, nipples and rings before use.

13-fl-oz can

- Rinse can lid; shake very well; open with a clean punch-type opener.
- Pour equal amounts of formula and water into bottle.
- Attach nipple; swirl gently to mix.
- Once feeding begins, **use within 1 hour or discard.**

1-QT bottle

- Shake very well before each use.
- Remove protective seal around bottle cap; twist off and clean cap. **Do not use if protective seal around cap or inner foil seal is damaged.**
- Invert cap; press down to pierce foil, then turn cap a half turn.
- Remove foil.
- Pour equal amounts of formula and water into bottle.
- Attach nipple; swirl gently to mix.
- Once feeding begins, **use within 1 hour or discard.**
- Prepare and feed all formula from this container **within 4 days after opening.**

Ready to Feed

Your baby's health depends on carefully following these directions. Ask your baby's doctor if you need to boil (sterilize) bottle, nipple and ring before use.

2-fl-oz bottle

- Do not add water.
- Shake very well.
- **Do not use if breakaway ring is missing or broken.**
- Twist off cap. Use clean hands to twist on nipple and ring (not included).
- Once feeding begins, **use within 1 hour or discard.**

8-fl-oz On-the-go bottle

- Do not add water.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Sensitive®

Infant Formula with Iron

- Shake very well.
- **Do not use if breakaway ring is missing or broken.**
- Twist off cap.
- Use clean hands to twist on clean nipple and ring (not included).
- For use in a different feeding container (cup or bottle):
 - Ask your baby's doctor if you need to boil (sterilize) the empty bottles before adding formula.
 - Pour desired amount into clean feeding container.
 - Recap bottle, refrigerate and **use within 48 hours.**
- No need to warm formula before feeding.
- Once feeding begins, **use within 1 hour or discard.**

1-QT bottle

- Do not add water.
- Shake very well before each use.
- Remove protective band, twist off and clean cap. **Do not use if band around cap or inner foil seal is damaged.**
- Invert cap; press down to pierce foil, then turn cap a half turn.
- Remove foil.
- Pour formula into bottle; attach nipple.
- Once feeding begins, **use within 1 hour or discard.**
- Prepare and feed all formula from the container **within 3 days after opening.**

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Storage & Handling

Powder

- Once mixed, store bottles in refrigerator and **feed to baby within 24 hours**.
- Store unopened or opened container at room temperature; avoid extreme temperatures.
- Use opened can contents **within 1 month**.
- Do not reuse container.

Concentrated Liquid

13-fl-oz can

- Cover opened cans and prepared formula not used immediately.
- Store in refrigerator and **feed to baby within 48 hours**.
- Store unopened cans at room temperature; avoid extreme temperatures.

1-QT bottle

- Once opened, store quart container in refrigerator.
- Store prepared bottles in refrigerator and **feed to baby within 48 hours**.
- Store unopened containers at room temperature; avoid extreme temperatures.
- Do not reuse container.
- Do not use this container to warm formula.

Ready to Feed

2-fl-oz-bottle

- Avoid prolonged exposure of bottles to light.
- Store unopened at room temperature; avoid extreme temperatures.

8-fl-oz On-the-go bottle

- Store unopened at room temperature; avoid extreme temperatures.
- Do not reuse bottle.
- **Protect from light. Store in carton until ready for use.**

1-QT bottle

- Once opened, store quart bottle in refrigerator.
- Store prepared bottles in refrigerator and **feed to baby within 48 hours**.
- Store unopened at room temperature; avoid extreme temperatures.
- Do not reuse bottle.
- Do not use this container to warm formula.

Product Information: Similac Total Comfort™

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Total Comfort™

Partially hydrolyzed protein infant formula with iron

- For discomfort due to persistent feeding issues – Similac Total Comfort has partially broken down protein for easy digestion.
- Our exclusive formula has:
 - Partially hydrolyzed protein for easy digestion.
 - Unique dual-carbohydrate blend for digestion and absorption.
 - Lutein and DHA for developing eyes and brain.
 - Calcium for strong bones - no palm olein oil.
 - Nucleotides to help support the immune system.
 - Prebiotics to help promote digestive health.
- Clinically shown to support normal infant growth.
- Suitable for infants with lactose sensitivity.
- Halal.
- Kosher.



Safety Precautions

- **Never use a microwave oven to warm formula.** Serious burns can result.
- Consult physician prior to switching formulas.
- Not for infants or children with galactosemia.
- **Warning:** Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

Ingredients

Powder Unflavored:

Corn Maltodextrin, Whey Protein Hydrolysate, High Oleic Safflower Oil, Sugar, Soy Oil, Coconut Oil, Galactooligosaccharides. Less than 2% of the Following: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Lycopene, Calcium Citrate, Potassium Phosphate, Calcium Phosphate, Potassium Citrate, Salt, Magnesium Chloride, Ascorbic Acid, Choline Chloride, Sodium Citrate, Calcium Hydroxide, Potassium Chloride, Ferrous Sulfate, Choline Bitartrate, Taurine, m-Inositol, Zinc Sulfate, Ascorbyl Palmitate, L-Carnitine, Carageenan, Niacinamide, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Potassium Iodide, Biotin, Sodium Selenate, Vitamin D3, Cyanocobalamin, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).

Allergens: Contains milk ingredients.

Availability

List Number	Item
62595	Similac Total Comfort Ready to Feed Institutional / 2-fl-oz (59-mL) bottles / 48 ct
62597	Similac Total Comfort Powder / 1.41-LB (638-g) SimplePac / 6 ct
62599	Similac Total Comfort Powder / 12.6-oz (357-g) can / 6 ct
63012	Similac Total Comfort Powder / 1.41-LB (638-g) SimplePac / 4 ct

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Similac Total Comfort™

Partially hydrolyzed protein infant formula with iron

Nutrition Information - Powder Unflavored

	100 Cal	1000 mL
	Value	Value
Osmolality, mOsm/kg H ₂ O	195	195
Protein, % Cal	9	9
Fat, % Cal	49	49
Carbohydrate, % Cal	42	42
Calories	100	676
Volume, mL	148	1000
Protein, g	2.32	15.68
Fat, g	5.4	36.5
Carbohydrate, g	11	74
Water, g	133	899
Linoleic Acid, mg	1000	6757
Vitamin A, IU	300	2027
Vitamin D, IU	60	405
Vitamin E, IU	1.5	10
Vitamin K, mcg	8	54
Thiamin (Vitamin B1), mcg	100	676
Riboflavin (Vitamin B2), mcg	150	1014
Vitamin B6, mcg	60	405
Vitamin B12, mcg	0.25	1.7
Niacin, mcg	1050	7095
Folic Acid (Folacin), mcg	15	101
Pantothenic Acid, mcg	450	3043
Biotin, mcg	4.4	29.8
Vitamin C (Ascorbic Acid), mg	9	61
Choline, mg	16	108
Inositol, mg	4.3	29
Calcium, mg	105	709
Phosphorus, mg	75	507
Magnesium, mg	6	40
Iron, mg	1.8	12
Zinc, mg	0.75	5.1
Manganese, mcg	5	34
Copper, mcg	90	609
Iodine, mcg	9	61
Selenium, mcg	1.8	12.2
Sodium, mg	44	297
Potassium, mg	118	797
Chloride, mg	65	439

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Preparation

Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- Wash your hands, surfaces and utensils.
- Pour water into clean bottle (see mixing guide).
- Add 1 unpacked level scoop (8.9 g) to each 2 fl oz of water.
- Return dry scoop to container.
- Cap bottle; shake well; attach nipple.
- Once feeding begins, **use within 1 hour or discard.**

Powder Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.9 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz
8 fl oz		4 scoops		8 fl oz

Pour prepared formula into feeding bottle, cap and store in refrigerator. **Use within 24 hours.**

Ready to Feed

Ready to feed formula if mixed with water may not provide proper nutrition, and repeated use of such feedings could cause illness.

- **Do not use if breakaway ring is missing or broken.**
- **Do not add water.**
- **Shake very well.**
- Twist off cap.
- Twist on nipple and ring. Use clean hands to attach nipple and ring.
- Once feeding begins, **use within 1 hour or discard.**

Storage & Handling

Powder

- Once mixed, store bottles in refrigerator and **feed to baby within 24 hours.**
- Store unopened or opened container at room temperature; avoid extreme temperatures.
- **Use opened can contents within 1 month.**
- Do not reuse container.

Ready to Feed

- Avoid prolonged exposure of bottles to light.
- Store unopened containers at room temperature; avoid extreme temperatures.
- Do not reuse bottle.

Product Information: Similac® 10-pct Glucose Water

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® 10-pct Glucose Water

Ready To Feed

- For initial or supplemental feeding as directed by physician.
- For oral use only.
- Use under medical supervision.
- Kosher.
- Halal.

Safety Precautions

- **Never use a microwave to warm product.** Serious burns can result.



Ingredients

Liquid Unflavored:

Water, and Dextrose Anhydrous.

Availability

List Number	Item
51004	Similac 10% Glucose Water Institutional / 2-fl oz (59-ml) bottle / 48 ct

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Similac® 10-pct Glucose Water

Ready To Feed

Nutrition Information - Liquid Unflavored

	1 fl oz	100 mL
	Value	Value
Water, %	93.7	93.7
Dextrose, %	10.0	10.0
Calories	12	40

Preparation

Directions for Preparation and Use

- Do not use if breakaway ring is missing or broken.
- Twist off cap.
- Twist on nipple and ring. Use clean hands to twist on nipple and ring.
- Once feeding begins, **use within one hour or discard.**

Storage & Handling

Directions for Use

- Store unopened at room temperature; avoid extreme temperatures.
- **Do not use bottles for breast milk storage.**

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Product Information: Similac® 5-pct Glucose Water

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® 5-pct Glucose Water

Ready To Feed

- For initial or supplemental feeding as directed by physician.
- For oral use only.
- Use under medical supervision.
- Kosher.
- Halal.

Safety Precautions

- **Never use a microwave to warm product.** Serious burns can result.



Ingredients

Liquid Unflavored:

Water, and Dextrose Anhydrous.

Availability

List Number	Item
51002	Similac 5% Glucose Water Institutional / 2-fl oz (59-mL) bottle / 48 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® 5-pct Glucose Water

Ready To Feed

Nutrition Information - Liquid Unflavored

	1 fl oz	100 mL
	Value	Value
Water, %	96.6	96.6
Dextrose, %	5.0	5.0
Calories	6	20

Preparation

Directions for Preparation and Use

- Do not use if breakaway ring is missing or broken.
- Twist off cap.
- Twist on nipple and ring. Use clean hands to twist on nipple and ring.
- Once feeding begins, **use within 1 hour or discard.**

Storage & Handling

Directions for Use

- Store unopened at room temperature; avoid extreme temperatures.
- **Do not use bottles for breast milk storage.**

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Product Information: Similac® Advance®

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® Advance®

Infant Formula with Iron

- A nutritionally complete, milk-based, iron-fortified infant formula for use as a supplement or alternative to breastfeeding.
- Our exclusive formula has:
 - Lutein and DHA for baby's developing eyes and brain.
 - Calcium for strong bones - no palm olein oil.
 - Nucleotides to help support the immune system.
 - Prebiotics to help promote digestive health.
 - Carotenoids naturally found in breast milk.
- Clinically proven tolerance in newborns and proven immune outcomes.
- Similac Advance has been clinically shown to support greater calcium absorption* for strong bones.
- Gluten-free.
- Kosher.
- Halal.



Safety Precautions

- **Never use a microwave oven to warm formula.** Serious burns can result.
- **Warning:** Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

* Similac® with Iron vs Enfamil® with Iron. Enfamil is not a trademark of Abbott Laboratories.

Ingredients

Powder Unflavored:

Nonfat Milk, Lactose, Whey Protein Concentrate, High Oleic Safflower Oil, Soy Oil, Coconut Oil, Galactooligosaccharides. Less than 2% of the Following: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Lycopene, Potassium Citrate, Calcium Carbonate, Ascorbic Acid, Soy Lecithin, Potassium Chloride, Magnesium Chloride, Ferrous Sulfate, Choline Bitartrate, Choline Chloride, Ascorbyl Palmitate, Salt, Taurine, m-Inositol, Zinc Sulfate, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, L-Carnitine, Vitamin A Palmitate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Cyanocobalamin, Calcium Phosphate, Potassium Phosphate, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).

Allergens: Contains milk and soy ingredients.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Availability

List Number	Item
53359	Similac Advance Powder / 1.45-lb (658-g) SimplePac / 6 ct
53363	Similac Advance Ready to Feed / 1-QT (946-mL) bottle / 6 ct
53365	Similac Advance Ready to Feed Institutional / 2-fl-oz (59-mL) bottle / 48 ct
53496	Similac Advance Powder / 2.13-lb (964-g) SimplePac / 6 ct
53498	Similac Advance Ready to Feed / 1-QT (946-mL) bottle / 6 ct
53526	Similac Advance On-the-go Ready to Feed / 8-fl-oz (237-mL) bottle / 12 x 1 ct
55957	Similac Advance Powder / 12.4-oz (352-g) can / 6 ct
56168	Similac Advance Ready to Feed / 1-QT (946-mL) bottle / 8 ct
56973	Similac Advance Concentrated Liquid / 13-fl-oz (384-mL) can / 12 ct
57600	Similac Advance Newborn Bottles Ready to Feed / 2-fl-oz (59 mL) bottle / 6 x 8 ct
57938	Similac Advance On-the-go Powder Singles / 0.61-oz (17.4-g) packet / 6 x 16 ct
58585	Similac Advance Powder / 1.93-lb (873-g) can / 6 ct
58605	Similac Advance On-the-go Ready to Feed / 8-fl-oz (237-mL) bottle / 4 x 6 ct
59952	Similac Advance On-the-go Ready to Feed / 8-fl-oz (237-mL) bottle / 24 ct
62811	Similac Advance On-the-go Powder Singles / 0.61-oz (17.4-g) packet / 4 x 16 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Powder Unflavored

	100 Cal*	1000 mL*
	Value	Value
Calories	100	676
Volume, mL	148	1000
Protein, g	2.07	14.00
Fat, g	5.60	37.8
Carbohydrate, g	10.71	72.4
Water, g	133	899
Linoleic Acid, mg	1000	6757
Potential Renal Solute Load, mOsm	18.7	126.7
Vitamin A, IU	300	2029
Vitamin D, IU	75	507
Vitamin E, IU	1.5	10.1
Vitamin K, mcg	8	54
Thiamin (Vitamin B1), mcg	100	676
Riboflavin (Vitamin B2), mcg	150	1014
Vitamin B6, mcg	60	406
Vitamin B12, mcg	0.25	1.69
Niacin, mcg	1050	7101
Folic Acid (Folacin), mcg	15	101
Pantothenic Acid, mcg	450	3043
Biotin, mcg	4.4	29.8
Vitamin C (Ascorbic Acid), mg	9	61
Choline, mg	16	108
Inositol, mg	4.7	31.8
Calcium, mg	78	528
Calcium, mEq	3.9	26.3
Phosphorus, mg	42	284
Magnesium, mg	6	41
Iron, mg	1.8	12
Zinc, mg	0.75	5.07
Manganese, mcg	5	34
Copper, mcg	90	609
Iodine, mcg	6	41
Selenium, mcg	1.8	12.2
Sodium, mg	24	162
Sodium, mEq	1.04	7.1
Potassium, mg	105	710
Potassium, mEq	2.7	18.2
Chloride, mg	65	440
Chloride, mEq	1.8	12.4

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Powder Unflavored Footnotes & References

Per 100 Cal*

*Nutrient values are applicable when prepared as directed.

Per 1000 mL*

*Nutrient values are applicable when prepared as directed.

Preparation

Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- Wash your hands, surfaces and utensils.

0.61-oz powder packet

- Pour **4 fl oz** of water into clean bottle.
- Slowly add one powder single (17.4 g) to bottle.
- Cap bottle; shake very well; attach nipple and ring.
- Once feeding begins, **use within 1 hour or discard.**

12.4-oz can, 1.45-LB, 1.93-LB & 2.13-LB SimplePac containers

- Pour water into clean bottle (see mixing guide).
- Add 1 unpacked level scoop (8.7 g) to each 2 fl oz of water.
- Return dry scoop to container.
- Cap bottle; shake well; attach nipple.
- Once feeding begins, **use within 1 hour or discard.**

Powder Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.7 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz
8 fl oz		4 scoops		8 fl oz

Concentrated Liquid

Your baby's health depends on carefully following these directions. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

13-fl-oz can

- Rinse can lid; shake very well; open with a clean punch-type opener.
- Pour equal amounts of formula and water into bottle.
- Attach nipple; swirl gently to mix.
- Once feeding begins, **use within 1 hour or discard.**

1-QT bottle

- Shake very well before each use.

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

- Remove protective seal around bottle cap; twist off and clean cap. **Do not use if band around cap or inner foil seal is damaged.**
- Invert cap; press down to pierce foil, then turn cap a half turn.
- Remove foil.
- Pour equal amounts of formula and water into bottle.
- Attach nipple; swirl gently to mix.
- Once feeding begins, **use within 1 hour or discard.**
- **Prepare and feed all formula from this container within 4 days after opening.**

Ready to Feed

Your baby's health depends on carefully following these directions. Ask your baby's doctor if you need to boil (sterilize) bottles, nipples and rings before use.

2-fl-oz bottle

- **Do not use if breakaway ring is missing or broken.**
- Do not add water.
- Shake very well.
- Twist off cap.
- Twist on nipple and ring.
- No need to warm before feeding.
- Once feeding begins, **use within 1 hour or discard.**

8-fl-oz On-the-go bottle

- **Do not use if breakaway ring is missing or broken.**
- Do not add water.
- Shake very well.
- Twist off cap.
- Twist on clean nipple and ring.
- No need to warm before feeding.
- Once feeding begins, **use within 1 hour or discard.**
- For use in a different feeding container (cup or bottle):
 - Ask your baby's doctor if you need to boil (sterilize) the empty bottles before adding formula.
 - Pour desired amount into clean feeding container.
 - Recap bottle, refrigerate and **use within 48 hours.**

1-QT bottle

- Do not add water.
- Shake very well before each use.
- Remove protective band, twist off and clean cap. **Do not use if band around cap or inner foil seal is damaged.**
- Invert cap; press down to pierce foil, then turn cap a half turn.
- Remove foil.
- Pour formula into bottle; attach nipple.
- Once feeding begins, **use within 1 hour or discard**
- Prepare and feed all formula from the 1-QT container **within 3 days after opening.**

Ready to Feed

Your baby's health depends on carefully following these directions. Ask your baby's doctor if you need to boil (sterilize) bottles, nipples and rings before use.

2-fl-oz bottle

- **Do not use if breakaway ring is missing or broken.**
- Do not add water.
- Shake very well.
- Twist off cap.
- Twist on nipple and ring.
- No need to warm before feeding.
- Once feeding begins, **use within 1 hour or discard.**

8-fl-oz On-the-go bottle

- **Do not use if breakaway ring is missing or broken.**
- Do not add water.
- Shake very well.
- Twist off cap.
- Twist on clean nipple and ring.
- No need to warm before feeding.
- Once feeding begins, **use within 1 hour or discard.**
- For use in a different feeding container (cup or bottle):
 - Ask your baby's doctor if you need to boil (sterilize) the empty bottles before adding formula.
 - Pour desired amount into clean feeding container.
 - Recap bottle, refrigerate and **use within 48 hours.**

1-QT bottle

- Do not add water.
- Shake very well before each use.
- Remove protective band, twist off and clean cap. **Do not use if band around cap or inner foil seal is damaged.**
- Invert cap; press down to pierce foil, then turn cap a half turn.
- Remove foil.
- Pour formula into bottle; attach nipple.
- Once feeding begins, **use within 1 hour or discard**
- Prepare and feed all formula from the 1-QT container **within 3 days after opening.**

Storage & Handling

Powder

0.61-oz powder packet

- Once mixed, feed immediately or cover and refrigerate; **use within 24 hours.**
- Store unopened packets at room temperature; avoid extreme temperatures.

12.4-oz can, 1.45-LB, 1.93-LB & 2.13-LB SimplePac containers

- Once mixed, store bottles in refrigerator and **feed to baby within 24 hours.**
- Store unopened or opened container at room temperature; avoid extreme temperatures.
- **Use opened can contents within 1 month.**
- Do not reuse container.

Concentrated Liquid

13-fl-oz can

- Cover opened cans and prepared formula not used immediately. Store in refrigerator and **feed to baby within 48 hours.**
- Store unopened cans at room temperature; avoid extreme temperatures.

1-QT bottle

- Once opened, store quart bottle in refrigerator.
- Store prepared bottles in refrigerator and **feed to baby within 48 hours.**
- Store unopened container at room temperature; avoid extreme temperatures.
- Do not reuse container.

Ready to Feed

2-fl-oz bottle & 8-fl-oz bottle

- Store unopened containers at room temperature; avoid extreme temperatures.
- Do not reuse bottle.
- **Protect from light.**
- **Store in carton until ready to use.**

1-QT bottle

- Once opened, store quart bottle in refrigerator
- Store prepared bottles in refrigerator and **feed to baby within 48 hours.**
- Store unopened containers at room temperature; avoid extreme temperatures.
- Do not reuse container.

Product Information: Similac® Advance® Organic

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® Advance® Organic

Infant Formula with Iron

- A nutritionally complete, organic, milk-based, iron-fortified infant formula for use as a supplement or alternative to breastfeeding.
- Our exclusive formula has:
 - Lutein and DHA for baby's developing eyes and brain.
 - Calcium for strong bones - no palm olein oil.
 - Nucleotides to help support the immune system.
 - Prebiotics to help promote digestive health.
 - Carotenoids naturally found in breast milk.
- Unique blend of organic nonfat milk, organic maltodextrin, and organic sugar.
- Certified USDA Organic.
- Kosher.
- Halal.
- Gluten-free.



Safety Precautions

- Never use a microwave oven to warm formula. Serious burns can result.
- **Warning:** Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

Ingredients

Powder Unflavored:

Organic Nonfat Milk, Organic Maltodextrin, Organic Sugar, Organic High Oleic Sunflower Oil, Organic Soy Oil, Organic Coconut Oil. Less than 2% of the Following: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Lycopene, Fructooligosaccharides, Potassium Citrate, Calcium Carbonate, Ascorbic Acid, Ascorbyl Palmitate, Ferrous Sulfate, Salt, Choline Chloride, Choline Bitartrate, Taurine, m-Inositol, Magnesium Chloride, Zinc Sulfate, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, L-Carnitine, Vitamin A Palmitate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Cyanocobalamin, Potassium Iodide, Potassium Hydroxide, and Nucleotides (Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate, Adenosine 5'-Monophosphate).

Allergens: Contains milk ingredients.

Availability

List Number	Item
50815	Similac Advance Organic Powder / 2.12-LB (936-g) SimplePac / 6 ct
50821	Similac Advance Organic Powder / 1.45-LB (657-g) SimplePac / 6 ct
56525	Similac Advance Organic Ready to Feed Institutional / 2-fl-oz (59-mL) Bottle / 48 ct
59883	Similac Advance Organic Ready to Feed / 1-QT (946-mL) Bottle / 6 ct
59885	Similac Advance Organic Ready to Feed / 1-QT (946-mL) Bottle / 6 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Powder Unflavored

	100 Cal*	1000 mL*
	Value	Value
Calories	100	676
Volume, mL	148	1000
Protein, g	2.07	14.07
Fat, g	5.63	38.26
Carbohydrate, g	10.4	70.7
Water, g	133	899
Linoleic Acid, mg	860	5844
Vitamin A, IU	300	2038
Vitamin D, IU	60	408
Vitamin E, IU	1.5	10.2
Vitamin K, mcg	8	54
Thiamin (Vitamin B1), mcg	100	679
Riboflavin (Vitamin B2), mcg	150	1019
Vitamin B6, mcg	60	408
Vitamin B12, mcg	0.25	1.70
Calcium, mg	78	530
Calcium, mEq	3.9	26.3
Niacin, mcg	1050	7135
Folic Acid (Folacin), mcg	15	102
Pantothenic Acid, mcg	450	3058
Biotin, mcg	4.4	29.9
Vitamin C (Ascorbic Acid), mg	9	61
Choline, mg	16	109
Inositol, mg	4.7	32
Phosphorus, mg	42	285
Magnesium, mg	6	41
Iron, mg	1.8	12.2
Zinc, mg	0.75	5.10
Manganese, mcg	5	34
Copper, mcg	90	612
Iodine, mcg	6	41
Selenium, mcg	1.8	12.2
Sodium, mg	24	163
Sodium, mEq	1.04	7.1
Potassium, mg	105	713
Potassium, mEq	2.7	18.1
Chloride, mg	65	442
Chloride, mEq	1.8	12.4

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Powder Unflavored Footnotes & References

Per 100 Cal*

*Nutrient values are applicable when prepared as directed.

Per 1000 mL*

*Nutrient values are applicable when prepared as directed.

Preparation

Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- Wash your hands, surfaces and utensils.
- Pour water into clean bottle (see mixing guide).
- Add 1 unpacked level scoop (8.6 g) to each 2 fl oz of water.
- Return dry scoop to holder in lid.
- Cap bottle; shake well; attach nipple.
- Once feeding begins, **use within 1 hour or discard.**

Powder Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.6 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz
8 fl oz		4 scoops		8 fl oz

Ready to Feed

Your baby's health depends on carefully following these directions. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to boil (sterilize) bottle, nipple and ring before use.

2-fl-oz bottle

- **Do not use if breakaway ring is missing or broken.**
- Do not add water.
 - Ready to feed formula if mixed with water may not provide proper nutrition, and repeated use of such feedings could cause illness.
- Shake very well.
- Twist off cap.
- Twist on nipple and ring.
 - Use clean hands to attach nipple and ring.
- Once feeding begins, **use within 1 hour or discard.**

1-QT bottle

- **Do not use if band around cap or inner foil seal is damaged.**
- Do not add water.
- Shake very well before each use.
- Remove protective band, twist off and clean cap.
- Invert cap; press down to pierce foil, then turn cap a half turn.
- Remove foil.
- Pour formula into bottle; attach nipple.
- Once feeding begins, **use within 1 hour or discard.**
- Prepare and feed all formula from the 1 QT container **within 3 days after opening.**

Storage & Handling

Ready to Feed

2-fl-oz bottle

- Avoid prolonged exposure of bottles to light.
- Store unopened at room temperature; avoid extreme temperatures.

1-QT bottle

- Once opened, store quart bottle in refrigerator.
- Store prepared bottles in refrigerator and **feed to baby within 48 hours**.
- Store unopened containers at room temperature; avoid extreme temperatures.
- Do not reuse container.

Powder

- Once mixed, store bottles in refrigerator and **feed to baby within 24 hours**.
- Store unopened or opened container at room temperature; avoid extreme temperatures.
- **Use opened container contents within 1 month.**
- Do not reuse container.

Product Information: Similac® For Supplementation

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® For Supplementation

Infant Formula with Iron

- For breastfeeding moms who choose to introduce formula.
- For a gentle introduction to formula.
- Gluten-free.
- Kosher.
- Halal.

Safety Precautions

- **Never use a microwave to warm product.** Serious burns can result.
- **Warning:** Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.



Ingredients

Powder Unflavored:

Nonfat Milk, Lactose, High Oleic Safflower Oil, Soy Oil, Coconut Oil, Galactooligosaccharides, Whey Protein Concentrate. Less than 2% of the Following: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Lycopene, Potassium Citrate, Calcium Carbonate, Ascorbic Acid, Soy Lecithin, Potassium Chloride, Magnesium Chloride, Ferrous Sulfate, Choline Bitartrate, Choline Chloride, Ascorbyl Palmitate, Salt, Taurine, m-Inositol, Zinc Sulfate, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, L-Carnitine, Vitamin A Palmitate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Cyanocobalamin, Calcium Phosphate, Potassium Phosphate, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
62990	Similac for Supplementation Ready to Feed / 2-fl-oz (59-mL) bottle / 6 x 8 ct
63013	Similac for Supplementation Powder / 1.45-LB (658-g) SimplePac / 4 ct
63092	Similac for Supplementation Ready to Feed Institutional / 2-fl-oz (59-mL) bottle / 48 ct

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Similac® For Supplementation

Infant Formula with Iron

Nutrition Information - Powder Unflavored

	100 Cal*	1000 mL*
	Value	Value
Calories	100	676
Volume, mL	148	1000
Protein, g	2.07	14.00
Fat, g	5.60	37.8
Carbohydrate, g	10.8	73.0
Water, g	133	899
Linoleic Acid, mg	1000	6757
Vitamin A, IU	300	2029
Vitamin D, IU	75	507
Vitamin E, IU	1.5	10.1
Vitamin K, mcg	8	54
Thiamin (Vitamin B1), mcg	100	676
Riboflavin (Vitamin B2), mcg	150	1014
Vitamin B6, mcg	60	406
Vitamin B12, mcg	0.25	1.69
Niacin, mcg	1050	7101
Folic Acid (Folacin), mcg	15	101
Pantothenic Acid, mcg	450	3043
Biotin, mcg	4.4	29.8
Vitamin C (Ascorbic Acid), mg	9	61
Choline, mg	16	108
Inositol, mg	4.7	31.8
Calcium, mg	78	528
Phosphorus, mg	42	284
Magnesium, mg	6	41
Iron, mg	1.8	12
Zinc, mg	0.75	5.07
Manganese, mcg	5	34
Copper, mcg	90	609
Iodine, mcg	6	41
Selenium, mcg	1.8	12.2
Sodium, mg	24	162
Potassium, mg	105	710
Chloride, mg	65	440

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® For Supplementation

Infant Formula with Iron

Powder Unflavored Footnotes & References

Per 100 Cal*

*Nutrient values are applicable when prepared as directed.

Per 1000 mL*

*Nutrient values are applicable when prepared as directed.

Preparation

Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottle, nipple and ring before use.

- Wash your hands, surfaces and utensils.
- Pour water into clean bottle (see mixing guide).
- Add 1 unpacked level scoop (8.7 g) to each 2 fl oz of water.
- Return dry scoop to container.
- Cap bottle; shake well; attach nipple.
- Once feeding begins, **use within 1 hour or discard.**

Powder Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.7 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz
8 fl oz		4 scoops		8 fl oz

Ready to Feed

Your baby's health depends on carefully following these. Ask your baby's doctor if you need to boil (sterilize) nipples and rings before use.

- Do not add water.
- Shake very well.
- **Do not use if breakaway ring is missing or broken.**
- Twist off cap.
- Twist on nipple and ring.
 - Since this formula is commercially sterile, use clean hands to attach any standard nipple and ring.
- No need to warm before feeding.
- **Once feeding begins, use within one hour or discard.**

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Storage & Handling

Powder

- Once mixed, store bottles in refrigerator and **feed to baby within 24 hours.**
- Store unopened or opened container at room temperature; avoid extreme temperatures.
- Do not store opened powder singles.
- **Use opened can contents within 1 month.**
- **Do not reuse container.**

Ready to Feed

- Avoid prolonged exposure of bottles to light.
- Store unopened at room temperature; avoid extreme temperatures.
- Do not reuse bottle.
- **Protect from light.**
- Store in carton until ready for use.

Product Information: Similac® Human Milk Fortifier

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® Human Milk Fortifier

- Similac Human Milk Fortifier is intended for low-birth-weight infants as a nutritional supplement to add to preterm human milk.
- Use under medical supervision.
- Infants fed Similac Human Milk Fortifier demonstrated excellent weight, length and head circumference gains.¹
- Specially formulated to mix easily with human milk.
- Low iron level provides flexibility to add iron as needed.
- A peer-reviewed clinical study showed that Similac Human Milk Fortifier, a low iron fortifier, preserved the antibacterial activity of breast milk against *E. coli*, *Staph*, GBS and *E. sakazakii* [now *C. sakazakii*].
- Well tolerated.
- Small, convenient packaging designed for less spillage.
- Halal.
- Kosher.



Safety Precautions

- Add only to human milk - do not add water.
- Tolerance to enteral feedings should be confirmed by offering small volumes of unfortified human milk. Once enteral feeding is well established, Similac Human Milk Fortifier can be added to human milk (see Preparation).
- Human milk can be fortified with Similac Human Milk Fortifier until the low-birth-weight infant reaches a weight of approximately 3600 g (approximately 8 lb) or as directed by a doctor.
- **Warning:** Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by a physician. Please follow directions as specified.
Improper dilution may be harmful.

* *Escherichia coli*, *Staphylococcus*, *Group B Streptococcus*, and *Enterobacter sakazakii* (renamed *Cronobacter sakazakii*)

¹ Barrett-Reis B et al. *Pediatrics* 2000;106:581-588.

² Chan GM. *J Perinatol* 2003;23:620-623.

Ingredients

Powder Unflavored:

Nonfat Milk, Whey Protein Concentrate, Corn Syrup Solids, Medium-Chain Triglycerides, Calcium Phosphate, Potassium Citrate. Less than 2% of the Following: Soy Lecithin, Minerals (Magnesium Chloride, Sodium Citrate, Calcium Carbonate, Salt, Potassium Phosphate, Zinc Sulfate, Ferrous Sulfate, Cupric Sulfate, Manganese Sulfate, Sodium Selenate), Vitamins (Vitamin A Palmitate, Vitamin D3, d-Alpha-Tocopheryl Acetate, Phylloquinone, Thiamin Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Cyanocobalamin, Niacinamide, Folic Acid, Calcium Pantothenate, Biotin, Ascorbic Acid, m-Inositol), and Potassium Hydroxide.

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
54598	Similac Human Milk Fortifier Powder Institutional / 0.9-g Packet / 3 x 50 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Powder Unflavored

	4 Packets (3.6 g)	1 Packet (0.9 g)
	Value	Value
Protein, g	1.0	0.25
Fat, g	0.36	0.09
Carbohydrate, g	1.8	0.45
Calories	14	3.5
Potential Renal Solute Load, mOsm	11.2	2.8
Vitamin A, IU	620	155
Vitamin D, IU	120	30
Vitamin E, IU	3.2	0.8
Vitamin K, mcg	8.3	2.1
Thiamin (Vitamin B1), mcg	233	58.3
Riboflavin (Vitamin B2), mcg	417	104
Vitamin B6, mcg	211	53
Vitamin B12, mcg	0.64	0.16
Niacin, mg	3.57	0.89
Folic Acid (Folacin), mcg	23	5.75
Pantothenic Acid, mg	1.5	0.37
Biotin, mcg	26	6.5
Vitamin C (Ascorbic Acid), mg	25	6.3
Calcium, mg	117	29.25
Calcium, mEq	5.84	1.46
Phosphorus, mg	67	16.8
Phosphorus, mEq	2.16	0.54
Magnesium, mg	7.0	1.75
Iron, mg	0.35	0.08
Zinc, mg	1.00	0.25
Manganese, mcg	7.2	1.8
Copper, mcg	170	42.5
Sodium, mg	15	3.75
Sodium, mEq	0.65	0.16
Potassium, mg	63	15.75
Potassium, mEq	1.61	0.40
Chloride, mg	38	9.5
Chloride, mEq	1.07	0.27

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Preparation

Powder

This product is for premature and low-birth-weight infants. Proper hygiene, handling and storage are important when preparing fortified human milk. Failure to follow these directions could result in severe harm.

Additional Calories Desired	Human Milk	Similac Human Milk Fortifier
2 (Cal/fl oz)	50 mL	1 packet
4 (Cal/fl oz)	25 mL	1 packet

- Gently shake container of human milk. Pour desired quantity into clean feeding container.
- To open packet, tear top off completely along perforation.
- Add fortifier to human milk and gently shake.
 - Bottle Feeding: Discard one hour after feeding begins.
 - Tube Feeding: Limit hang time to a maximum of 4 hours.

Storage & Handling

Powder

- Use immediately or store unused fortified human milk in the refrigerator; use within 24 hours or according to hospital procedures for handling human milk.
- Gently shake before using.
- Store unopened packets at room temperature; avoid extreme temperatures.

Product Information: Similac® Human Milk Fortifier Concentrated Liquid

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® Human Milk Fortifier Concentrated Liquid

- Now available as a concentrated liquid. Similac Human Milk Fortifier delivers the same nutrient profile as the one healthcare professionals have been relying on for over a decade.*
- Small, convenient packet is designed for easy mixing.
- Similac Human Milk Fortifier Concentrated Liquid, when added to human milk, meets the nutrient recommendations for the premature infant.¹
- Similac Human Milk Fortifier Concentrated Liquid is commercially sterile and meets the ADA and CDC recommendation to use liquid for NICU feedings.^{† 2,3}
- Similac Human Milk Fortifier Concentrated Liquid packet is simple to open and mixes easier with human milk than powder.⁴
- Low iron level provides flexibility to add iron as needed.
- DHA and ARA - same as Similac Special Care Infant Formulas.[§]



Safety Precautions

- This product is nutritionally incomplete.
- **Never use a microwave oven to warm feedings.** Serious burns can result.
- Add only to human milk - do not add water.

* As fed when mixed with human milk.

† American Dietetic Association and Centers for Disease Control and Prevention

‡ Data on file, 2010. Abbott Nutrition Market Research, Abbott Laboratories, Columbus, Ohio.

§ When mixed with human milk.

¹ Klein CJ. Nutrient requirements for preterm infant formulas. *J Nutr* 2002;132(6):1395S-1577S.

² Centers for Disease Control and Prevention. *Enterobacter sakazakii* infections associated with the use of powdered infant formula—Tennessee, 2001. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5114a1.htm>. Accessed May 28, 2010.

³ Robbins ST, Beker LT. *Infant Feedings: Guidelines for Preparation of Formula and Breastmilk in Health Care Facilities*. American Dietetic Association, 2011.

⁴ Data on file, 2010. Abbott Nutrition Market Research, Abbott Laboratories, Columbus, Ohio.

Ingredients

Concentrated Liquid Unflavored:

Water, Nonfat Milk, Corn Syrup Solids, Medium-Chain Triglycerides, Whey Protein Concentrate. Less than 2% of the Following: Minerals (Calcium Phosphate, Magnesium Chloride, Potassium Citrate, Salt, Calcium Carbonate, Potassium Phosphate, Sodium Citrate, Magnesium Phosphate, Zinc Sulfate, Ferrous Sulfate, Cupric Sulfate, Manganese Sulfate and Sodium Selenate), Vitamins (Vitamin A Palmitate, Vitamin D3, d-Alpha-Tocopheryl Acetate, Phylloquinone, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Cyanocobalamin, Niacinamide, Folic Acid, Calcium Pantothenate, Biotin, Ascorbic Acid, m-Inositol and Choline Chloride), Soy Lecithin, M. Alpina Oil, C. Cohnii Oil, and Potassium Hydroxide.

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
56649	Similac Human Milk Fortifier Concentrated Liquid Instit / 5-mL (0.17-fl-oz) packet / 6 x 24 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® Human Milk Fortifier Concentrated Liquid

Nutrition Information - Concentrated Liquid Unflavored

	100 Cal Prepared*	5 mL packet	30 mL Prepared*
	Value	Value	Value
Volume, mL	127	5	30
Calories	100	6.85	24
Protein, g	2.97	0.349	0.701
Fat, g	5.24	0.266	1.24
Carbohydrate, g	10.4	0.807	2.47
Water, g	108	3.64	25.6
Linoleic Acid, mg	395 [†]	1.00	93.3 [†]
Vitamin A, IU	1248	197	294
Vitamin D, IU	150	34.9	35.4
Vitamin E, IU	5.26	0.973	1.24
Vitamin K, mcg	10.5	2.43	2.48
Thiamin (Vitamin B1), mcg	313	68.7	73.9
Riboflavin (Vitamin B2), mcg	573	123	135
Vitamin B6, mcg	279	62.0	65.7
Vitamin B12, mcg	0.400	0.0825	0.0943
Niacin, mcg	4586	1044	1082
Folic Acid (Folacin), mcg	32.4	6.82	7.65
Pantothenic Acid, mg	2073	444	489
Biotin, mcg	32.5 [†]	7.57	7.67 [†]
Vitamin C (Ascorbic Acid), mg	44.0	7.71	10.4
Choline, mg	13.8 [†]	0.919	3.27
Inositol, mg	22.9 [†]	1.72	5.41 [†]
Calcium, mg	175	35.1	41.3
Phosphorus, mg	98.3	20.0	23.2
Magnesium, mg	12.4	2.16	2.93
Iron, mg	0.582	0.107	0.137
Zinc, mg	1.65	0.304	0.390
Manganese, mcg	9.62 [†]	2.12	2.27 [†]
Copper, mcg	289	52.0	68.1
Iodine, mcg	13.3 [†]	0.470	3.15 [†]
Selenium, mcg	2.43 [†]	0.204	0.574 [†]
Sodium, mg	49.1	5.39	11.6
Potassium, mg	148	20.7	35.0
Chloride, mg	115	13.4	27.2

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® Human Milk Fortifier Concentrated Liquid

Concentrated Liquid Unflavored Footnotes & References

Per 100 Cal Prepared*

*Per 100 Calories, as fed, approximate 4 packets in 100 mL preterm human milk (24 Cal/fl oz). Preterm human milk values from "Meeting the Special Needs of Low-Birth-Weight and Premature Infants in the Hospital." Columbus, OH: Abbott Nutrition, Abbott Laboratories, Jan 1998, p 56.

†Values based on term human milk from Academy of Pediatrics Committee on Nutrition. "Pediatric Nutrition Handbook", ed 4. Elk Grove Village: American Academy of Pediatrics, 1998:40; 132-135, 217, 258, 655-658.

Per 30 mL Prepared*

*One packet (5 mL) Similac Human Milk Fortifier Concentrated Liquid in 25 mL preterm human milk. Preterm human milk values from "Meeting the Special Needs of Low-Birth-Weight and Premature Infants in the Hospital." Columbus, OH: Abbott Nutrition, Abbott Laboratories, Jan 1998, p 56.

†Values based on term human milk from Academy of Pediatrics Committee on Nutrition. "Pediatric Nutrition Handbook", ed 4. Elk Grove Village: American Academy of Pediatrics, 1998:40; 132-135, 217, 258, 655-658.

Preparation

Directions for Use:

This product is for premature and low-birth-weight infants. Proper hygiene, handling and storage are important when preparing fortified human milk. **Follow directions as specified. Improper dilution may be harmful.**

Additional Calories Desired	Human Milk	Similac Human Milk Fortifier	Approximate Yield
2 Calories/fl oz	50 mL	1 packet	55 mL
4 Calories/fl oz	25 mL	1 packet	30 mL

Immediate Use

- Pour desired quantity of human milk into sterile feeding container.
- Product separation is normal. Massage packet contents until uniform in color.
- To open packet, tear top off completely along perforation.
- Add contents to human milk and **shake until mixed well**. If bottle feeding, **use within one hour or discard**.

When Stored

- Divide unused fortified human milk into individual bottles or syringes. Refrigerate and use within 24 hours.
- After refrigeration, warm to room temperature.
- Settling may occur; **shake for 10 seconds or until mixed well**. Feed promptly.
- For syringes, allow 5 mL headspace for mixing. Feed vertically, tip down.

Storage & Handling

Directions for Storage:

- Keep unopened packets at room temperature in the protective carton; avoid extreme temperatures.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Product Information: Similac® Nipples

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

- Provides the flexibility to choose a ready-for-use nipple and ring that match each infant's sucking strength or need.
- **Infant** (latex-free)—for infants with developed sucking strength.
- **Premature** (latex-free)—for preterm infants with less developed sucking strength.
- **Orthodontic** (contains latex)—for newborn infants, to stimulate oral exercise while feeding.
- **Slow Flow** (latex-free)—single hole for a more controlled flow rate during feeding.
- **Cleft Palate Assembly** (latex-free)—uniquely designed to meet the feeding needs of the infant with cleft palate. Requires minimal sucking. Formula flow can be regulated by gently squeezing the side walls of the attached bottle.
- Nipples and rings individually prepackaged and ready for use.
- Nipples and rings form a seal to prevent leakage during feeding.
- Disposable—not for reuse.
- Easy to use.



Safety Precautions

- DO NOT use as a pacifier, as it presents a choking hazard.
- DO NOT reuse, as reuse can cause deterioration or separation of the rubber, presenting a choking hazard.
- Keep out of reach of children.

Availability

List Number	Item
00079	Similac Infant Nipple & Ring / 250 ct
00094	Similac Premature Nipple & Ring / 250 ct
50512	Similac Orthodontic Nipple & Ring / 250 ct
53894	Similac Slow Flow Nipple and Ring / 250 ct
56054	Cleft Palate Assembly / 3 ct
56366	Similac Slow Flow Nipple & Ring Kit / 50 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Product Information: Similac® PM 60/40

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® PM 60/40

Low-Iron Infant Formula

- For infants who would benefit from lowered mineral intake, including those with impaired renal function.
- Use under medical supervision.
- Mineral levels closely approximating the mineral content of human milk (60:40 ratio of whey to casein).
- Calcium-to-phosphorus ratio and content designed to treat serum calcium disorders—both hypercalcemia and hypocalcemia due to hyperphosphatemia.
- Gluten-free.
- Halal.
- Kosher.

Safety Precautions

- In conditions where the infant is losing abnormal quantities of one or more electrolytes, it may be necessary to supply electrolytes from sources other than the formula.
- Additional iron should be supplied from other sources.
- Never use a microwave to warm formula. Serious burns can result.
- Not for IV use.



Ingredients

Powder Unflavored:

Lactose, High Oleic Safflower Oil, Whey Protein Concentrate, Soy Oil, Coconut Oil, Sodium Caseinate. Less than 2% of the Following: Potassium Citrate, Calcium Phosphate, Calcium Carbonate, Magnesium Chloride, Potassium Chloride, Ascorbic Acid, m-Inositol, Salt, Choline Chloride, Taurine, Ascorbyl Palmitate, Ferrous Sulfate, Zinc Sulfate, Mixed Tocopherols, L-Carnitine, Niacinamide, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Beta-Carotene, Folic Acid, Manganese Sulfate, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Cyanocobalamin, and Potassium Hydroxide.

Allergens: Contains milk ingredients.

Availability

List Number	Item
00850	Similac PM 60/40 Powder / 14.1-oz (400-g) can / 6 ct

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Nutrition Information - Powder Unflavored

	100 Cal	1000 mL
	Value	Value
Osmolality, mOsm/kg H ₂ O	280	280
Calories	100	676
Volume, mL	148	1000
Protein, g	2.2	15
Fat, g	5.6	37.9
Carbohydrate, g	10.2	69.0
Water, g	134	899
Linoleic Acid, mg	1000	6763
Potential Renal Solute Load, mOsm	18.3	124.1
Vitamin A, IU	300	2029
Vitamin D, IU	60	406
Vitamin E, IU	1.5	10.1
Vitamin K, mcg	8	54
Thiamin (Vitamin B1), mcg	100	676
Riboflavin (Vitamin B2), mcg	150	1014
Vitamin B6, mcg	60	406
Vitamin B12, mcg	0.3	1.7
Niacin, mcg	1050	7101
Folic Acid (Folacin), mcg	15	101
Pantothenic Acid, mcg	450	3043
Biotin, mcg	5	30.4
Vitamin C (Ascorbic Acid), mg	9	61
Choline, mg	12	81
Inositol, mg	24	162
Calcium, mg	56	379
Calcium, mEq	2.8	18.9
Phosphorus, mg	28	189
Magnesium, mg	6	40.6
Iron, mg	0.7	4.7
Zinc, mg	0.8	5.1
Manganese, mcg	5	34
Copper, mcg	90	609
Iodine, mcg	6	41
Selenium, mcg	1.8	12.2
Sodium, mg	24	162
Potassium, mg	80	541
Chloride, mg	59	399

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Preparation

Directions for Preparation and Use

Your baby's health depends on carefully following these easy directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to boil (sterilize) bottles, nipples and rings before use.

- Wash your hands, surfaces and utensils.
- Pour desired amount of water into a clean bottle (see mixing guide).
- Add 1 unpacked level scoop (8.7 g) to each 2 fl oz of water
- Return dry scoop to can.
- Cap bottle; shake well; attach nipple.
- Once feeding begins, **use within 1 hour or discard.**

Powder Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.7 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz

For a larger amount of formula, add 1 unpacked, level cup of powder (100 g) to 23 fl oz of water (makes approximately 26 fl oz of formula).

When mixed as directed, the contents of this can will make approximately 102 fl oz of formula.

Storage & Handling

Powder

- Once mixed, store bottles in refrigerator and **feed to baby within 24 hours.**
- Store unopened or opened container at room temperature; avoid extreme temperatures.
- **Use opened can contents within 1 month.**

Product Information: Similac® Prenatal & Breastfeeding Dietary Supplement

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® Prenatal & Breastfeeding Dietary Supplement

- Similac Prenatal provides nutrients to support both mom and baby during pregnancy and breastfeeding*
- Meets 100% of the recommended Daily Value for 13 essential vitamins and minerals
- Similac Prenatal goes beyond the essentials: DHA and lutein in addition to 22 essential vitamins and minerals
- Similac Prenatal to help replenish important nutrients for mom during pregnancy and breastfeeding*
- Natural vitamin E to support immune health*
- Vitamin D is essential to help support calcium absorption for bone growth*
- Iron to help support fetal brain development*
- Folic acid to help support fetal brain development*



Safety Precautions

- **Warning:** Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. **Keep this product out of reach of children.** In case of accidental overdose, call a doctor or poison control center immediately.
- Always inform your doctor about the use of dietary supplements.
- Do not use this product in combination with other dietary supplements.

* **These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.**

* **These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.**

Ingredients

Multivit MinTablet Unflavored:

Other Ingredients: Microcrystalline Cellulose, Croscarmellose Sodium, Hypromellose, Titanium Dioxide, Polydextrose, Silicon Dioxide, Magnesium Stearate, Talc, Maltodextrin, and Medium-Chain Triglycerides.

Allergens: Manufactured in a facility that processes fish, milk, shellfish, tree nut and wheat ingredients.

Availability

List Number	Item
62427	Similac Prenatal Dietary Supplement / 4 x 3 30-ct each multivit/min tablets and DHA/lutein softgels

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® Prenatal & Breastfeeding Dietary Supplement

Nutrition Information - Multivit MinTablet Unflavored

	1 Tablet
	Value
Vitamin A (50% as Beta-Carotene, 50% Vitamin A Palmitate), IU	4000
Vitamin C (as Ascorbic Acid), mg	120
Vitamin D (as Cholecalciferol), IU	600
Thiamin (as Thiamine Mononitrate), mg	1.8
Riboflavin, mg	1.7
Niacin (as Niacinamide), mg	20
Vitamin B6 (as Pyridoxine Hydrochloride), mg	2.6
Folic Acid, mcg	800
Vitamin B12 (as Cyanocobalamin), mcg	8
Biotin, mcg	35
Pantothenic Acid (as Calcium D-Pantothenate), mg	10
Calcium (as Calcium Carbonate), mg	300
Iron (as Ferrous Fumarate), mg	27
Iodine (as Potassium Iodide), mcg	210
Magnesium (as Magnesium Oxide), mg	200
Zinc (as Zinc Oxide), mg	15
Selenium (as Sodium Selenate), mcg	70
Copper (as Copper Gluconate), mg	2
Manganese (as Manganese Gluconate), mg	2.6
Chromium (as Chromium Chloride), mcg	45
Molybdenum (as Sodium Molybdate), mcg	50

Multivit MinTablet Unflavored Footnotes & References

Per 1 Tablet

*% Daily Value for pregnant/lactating women.

Preparation

Directions

- Do not use this product if safety seal underneath bottle cap is torn or missing.
- During pregnancy and lactation, take **one tablet and one softgel** daily with food.
- Eating a healthy diet along with your recommended daily prenatal is important for both you and your baby.

Storage & Handling

- Store with cap secured at room temperature.
- Avoid extreme temperatures.

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Product Information: Similac® Simply Smart™

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

- Our experts partnered with moms to design the one bottle that has it all.
- Nipple designed for feeding between breast and bottle.
- SmartClose™ locks out leaks
- EasyMix™ is designed to help reduce bubbles and clumps.
- **Cautions**
 - Never use the microwave to warm formula. Serious burns can result.
 - It is very important to clean the bottle and all the components before and after every feeding.
 - Remove the bottle from your baby's mouth when not feeding. Prolonged or continuous fluid sucking can cause tooth decay.
 - The components of the bottle are not toys and should not be given to children to play with or suck on.
 - Prior to every use, inspect nipples. Replace the nipple if it shows any signs of wear, such as tearing, cracking or a change of shape.



Availability

List Number	Item
62016	Similac Simply Smart Bottle 4-oz Bottle single / 4 x 3 ct
62024	Similac Simply Smart Level 1 Nipple (Slow Flow) 2 count / 6 x 2 ct
62026	Similac Simply Smart Level 2 Nipple (Medium Flow) 2 count / 6 x 2 ct
62028	Similac Simply Smart Level 3 Nipple (Fast Flow) 2 count / 6 x 2 ct
62030	Similac Simply Smart Level Y Nipple (for Thicker Formula) 2 count / 6 x 2 ct
62032	Similac On-the-go Powder Cap single / 6 x 2 ct
62053	Similac Simply Smart 4-oz Bottle 2 count / 6 x 2 ct
62054	Similac Simply Smart Starter Set / 2 ct
62055	Similac Simply Smart 8-oz Bottle single / 4 x 3 ct
62057	Similac Simply Smart 8-oz Bottle 2 count / 6 x 2 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Preparation

Instructions for Use: Bottle & Nipples

Before first use

- Wash all bottle components in hot, soapy water before initial use. Ask your baby's doctor if you need to boil or sterilize bottles, nipples and rings.

Cleaning

- Wash bottle and all components before every use.
- Wash your hands, surfaces and utensils.
- Dishwasher safe; use top rack only and wash on a normal cycle.

Bottle assembly and feeding instructions

- Insert EasyMix™ into the top of the bottle for a secure fit. *Note: The bottle will function with or without the EasyMix™.*
- Put desired amount of breast milk or formula into the bottle. When using formula, mix according to the formula manufacturer's instructions.
- Insert the nipple by pulling the top of the nipple through the SmartClose™ ring until it fits securely.
- Tighten the ring on the bottle. When it is securely locked you will see blue in the indicator window.
- Shake or swirl the bottle. The EasyMix™ has been designed to help reduce the amount of bubbles or formula clumps.
- To test the temperature of the breast milk or formula, we recommend that you always sprinkle a few drops on the inside of your wrist to make sure it is not too hot.

Instructions for Use: On-the-go Powder Cap (sold separately)

- Wash and dry the powder cap before every use.
- Screw both parts of the cap together.
- Open the lid and fill with enough powder formula to make 1 bottle. Refer to the formula manufacturer's instructions for the appropriate amount of formula for your baby.
- Close the lid and attach this cap to bottle. Formula should not be stored in the cap for **longer than 5 days**.
- Add water to bottle as directed by formula manufacturer.
- Just prior to feeding, pour powder formula into bottle.
- After use, unscrew cap into two parts and clean with hot, soapy water. Allow the cap to thoroughly dry before using again by storing it in a cool, dry place.

Product Information: Similac® Soy Isomil®

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® Soy Isomil®

Soy Infant Formula with Iron

- Comfort your baby's fussiness and gas with Similac Soy Isomil — specially designed with the gentleness of soy to soothe the tummy.
- Complete nutrition for sensitive tummies.
- Soy for fussiness and gas.
- #1 formula for complete soy nutrition.

Safety Precautions

- Never use a microwave oven to warm mixture. Serious burns can result.
- **Warning:** Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.



Ingredients

Powder Unflavored:

Corn Syrup Solids (39%), Soy Protein Isolate (15%), High Oleic Safflower Oil (11%), Sugar (10%), Soy Oil (8%), Coconut Oil (8%). Less than 2% of the Following: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Lycopene, Fructooligosaccharides, Calcium Phosphate, Potassium Citrate, Potassium Chloride, Magnesium Chloride, Salt, Ascorbic Acid, Choline Chloride, L-Methionine, Taurine, Ascorbyl Palmitate, Ferrous Sulfate, m-Inositol, Mixed Tocopherols, Zinc Sulfate, d-Alpha-Tocopheryl Acetate, L-Carnitine, Niacinamide, Calcium Pantothenate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Potassium Hydroxide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, and Cyanocobalamin.

Allergens: Contains soy ingredients.

Availability

List Number	Item
50819	Similac Soy Isomil Powder / 1.45-lb (658-g) SimplePac / Case of 6
50825	Similac Soy Isomil Powder / 2.13-lb (964-g) SimplePac / Case of 6
55963	Similac Soy Isomil Powder / 12.4-oz (352-g) Can / Case of 6
55967	Similac Soy Isomil Ready to Feed / 1-QT (946-mL) Bottle / Case of 6
56308	Similac Soy Isomil Ready to Feed Institutional / 2-fl-oz (59-mL) Bottle / Case of 48
56975	Similac Soy Isomil Concentrated Liquid / 13-fl-oz (384-mL) Can / Case of 12
56982	Similac Soy Isomil On-the-go Powder Singles / 0.62-oz (17.6-g) Packets / Case of 96
58292	Similac Soy Isomil Ready to Feed 6-Pack / 1-QT (946-mL) Bottle
58601	Similac Soy Isomil On-the-Go / 8-fl-oz (237-mL) Bottle / Case of 24

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Nutrition Information - Powder Unflavored

	100 Cal*	1000 mL*
	Value	Value
Protein, % Cal	10	10
Protein Source	Soy Protein Isolate, L-Methionine	Soy Protein Isolate, L-Methionine
Fat, % Cal	49	49
Fat Source	High Oleic Safflower, Soy and Coconut Oils (0.15% DHA, 0.40% ARA)	High Oleic Safflower, Soy and Coconut Oils (0.15% DHA, 0.40% ARA)
Fat Ratio	41:30:28	41:30:28
Carbohydrate, % Cal	41	41
Carbohydrate Source	Corn Syrup Solids, Sugar, Fructooligosaccharides (FOS)	Corn Syrup Solids, Sugar, Fructooligosaccharides (FOS)
Prebiotic	FOS	FOS
Carbohydrate Ratio	78:19:3	78:19:3
Osmolality, mOsm/kg H2O	200	200
Calories	100	676
Volume, mL	148	1000
Protein, g	2.45	16.57
Fat, g	5.46	36.93
Carbohydrate, g	10.4	70.3
Water, g	133	899
Linoleic Acid, mg	1000	6763
Potential Renal Solute Load, mOsm	22.8	154.5
Vitamin A, IU	300	2029
Vitamin D, IU	60	406
Vitamin E, IU	1.5	10.1
Vitamin K, mcg	11	74
Thiamin (Vitamin B1), mcg	60	406
Riboflavin (Vitamin B2), mcg	90	609
Vitamin B6, mcg	60	406
Vitamin B12, mcg	0.45	3.04
Niacin, mcg	1350	9130
Folic Acid (Folacin), mcg	15	101
Pantothenic Acid, mcg	750	5072
Biotin, mcg	4.5	30.4
Vitamin C (Ascorbic Acid), mg	9	61
Choline, mg	12	81
Inositol, mg	5	33.8
Calcium, mg	105	710
Calcium, mEq	5.2	35.4
Phosphorus, mg	75	507
Magnesium, mg	7.5	50.7

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Nutrition Information - Powder Unflavored

	100 Cal [*]	1000 mL [*]
	Value	Value
Iron, mg	1.8	12.2
Zinc, mg	0.75	5.07
Manganese, mcg	25	169
Copper, mcg	75	507
Iodine, mcg	15	101
Selenium, mcg	1.8	12.2
Sodium, mg	44	298
Sodium, mEq	1.9	12.9
Potassium, mg	108	730
Potassium, mEq	2.8	18.7
Chloride, mg	62	419
Chloride, mEq	1.8	11.8

Powder Unflavored Footnotes & References

Per 100 Cal^{*}

^{*}Nutrient values are applicable when prepared as directed.

Per 1000 mL^{*}

^{*}Nutrient values are applicable when prepared as directed.

Preparation

Powder Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.8 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz
8 fl oz		4 scoops		8 fl oz

Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

0.62-oz powder packet

- Wash your hands, surfaces and utensils.
- Pour **4 fl oz** of water into clean bottle.
- Slowly add one powder single (17.6 g) to bottle.
- Cap bottle; shake very well; attach nipple and ring.
- Once feeding begins, **use within 1 hour or discard**.

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Similac® Soy Isomil®

Soy Infant Formula with Iron

12.4-oz can & 1.45-LB and 2.12-LB SimplePac containers

- Wash your hands, surfaces and utensils.
- Pour water into clean bottle (see mixing guide).
- Add 1 unpacked level scoop (8.8 g) to each 2 fl oz of water.
- Return dry scoop to container.
- Cap bottle; shake well; attach nipple.
- Once feeding begins, **use within 1 hour or discard.**

Powder Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.8 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz
8 fl oz		4 scoops		8 fl oz

Concentrated Liquid

Your baby's health depends on carefully following these directions. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- Rinse can lid; shake very well; open with a clean punch-type opener.
- Pour equal amounts of formula and water into bottle.
- Attach nipple; swirl gently to mix.
- Once feeding begins, **use within 1 hour or discard.**

Ready to Feed

Your baby's health depends on carefully following these directions. Ask your baby's doctor if you need to boil (sterilize) bottles, nipples and rings before use.

2-fl-oz bottle

- Do not use if breakaway ring is missing or broken.
- Do not add water.
- Shake very well.
- Twist off cap; use clean hands to twist on nipple and ring.
- Once feeding begins, **use within 1 hour or discard.**

8-fl-oz On-the-go bottle

- Do not use if breakaway ring is missing or broken.
- Do not add water.
- Shake very well.
- Twist-off cap.
- Twist-on clean nipple and ring (not included).
- For use in a **different feeding container** (cup or bottle):
 - Ask your baby's doctor if you need to boil (sterilize) the empty bottles before adding formula.
 - Pour desired amount into clean feeding container. Recap bottle, refrigerate and **use within 48 hours.**
- No need to warm formula before feeding.
- Once feeding begins, **use within 1 hour or discard.**

1-QT bottle

- Do not use if band around cap or inner foil seal is damaged.
- Do not add water.
- Shake very well before each use.
- Remove protective band, twist off and clean cap.
- Invert cap; press down to pierce foil, then turn cap a half turn.
- Remove foil.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® Soy Isomil®

Soy Infant Formula with Iron

- Pour formula into bottle; attach nipple.
- Once feeding begins, **use within 1 hour or discard.**
- Prepare and feed all formula from the 1 QT container **within 3 days after opening.**

Storage & Handling

Powder

0.62-oz powder packet

- Once mixed, feed immediately or cover and refrigerate; **use within 24 hours.**
- Store unopened packets at room temperature; avoid extreme temperatures.

12.4-oz can & 1.45-LB and 2.12-LB SimplePac containers

- Once mixed, store bottles in refrigerator and **feed to baby within 24 hours.**
- Store unopened or opened container at room temperature; avoid extreme temperatures.
- Use opened container contents **within 1 month.**
- Do not reuse container.

Concentrated Liquid

- Cover opened cans and prepared formula not used immediately.
- Store in refrigerator and **feed to baby within 48 hours.**
- Store unopened cans at room temperature; avoid extreme temperatures.

Ready to Feed

2-fl-oz bottle

- Avoid prolonged exposure of bottles to light.
- Store unopened at room temperature; avoid extreme temperatures.

8-fl-oz On-the-go bottle

- Store unopened containers at room temperature; avoid extreme temperatures.
- Do not reuse bottle.
- **Protect from light.**
- **Store in carton until ready to use.**

1-QT bottle

- Once opened, store quart bottle in refrigerator.
- Store bottles in refrigerator and **feed to baby within 48 hours.**
- Store unopened containers at room temperature; avoid extreme temperatures.
- Do not reuse container.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Product Information: Similac® Sterilized Water

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® Sterilized Water

Ready To Feed

- As an initial feeding.
- Use under medical supervision.
- Halal.
- Kosher.

Safety Precautions

- For oral use only.
- **Never use a microwave to warm product.** Serious burns can result.



Ingredients

Liquid Unflavored:
Water

Availability

List Number	Item
51000	Similac Water (Sterilized) Institutional / 2 fl oz (59 mL) Bottle / 48 ct
58037	Similac Water (Sterilized) Institutional / 1 QT 1.8 fl oz (1 L) Bottle / 8 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Liquid Unflavored

	1 fl oz
	Value
Water, fl oz	1

Preparation

Instructions for Use

2-fl-oz bottle

- Do not use if ring is missing or broken.
- Twist off cap.
- Twist on nipple and ring; use clean hands to attach nipple and ring.
- **Once feeding begins, use within 1 hour or discard.**

Storage & Handling

Instructions for Use

2-fl-oz bottle

- Store unopened at room temperature; avoid extreme temperatures.
- Do not use for breast milk storage.

1-L bottle

- After opening, replace cap, refrigerate and **use within 48 hours.**
- Store unopened bottles in a cool place; avoid excessive heat.
- Do not reuse bottle.

Product Information: Similac® Volu-Feed®

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® Volu-Feed®

Volumetric Feeding System

- When accurate measurement of formula intake is needed (for example, with premature and low-birth-weight infants).
- Easier to Read: even numbers on one side and odd numbers on the other side; numbers centered on the graduation marks; and larger, easier to read font.
- Easier to Write: Double the writing panel space.
- Lighter-weight plastic: 50,000 pounds less plastic per year (12.5% less plastic per bottle); polypropylene.
- Food-grade quality.
- DEHP- and Bisphenol A-free.
- Ready for Use—in a clean-peel blisterpack.
- Nipple ready.



Availability

List Number	Item
00081	Volu-Feed Dispensing Caps / 250 ct
00180	Volu-Feed Nurser / 100 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com