

# D08 Shaking hands in the face of danger? Interpersonal distance, peripersonal space, pro- and anti-social consequences of face masks during the COVID-19 pandemic

Katharina Kühne\* MSc., Martin H. Fischer PhD., Melinda A. Jeglinski-Mende MSc.  
University of Potsdam, Germany \*Corresponding author: Katharina Kühne [kkuehne@uni-potsdam.de](mailto:kkuehne@uni-potsdam.de)

## Introduction

- **Peripersonal space (PPS):** space surrounding our body, where multisensory integration of stimuli and action execution take place (Rizzolatti et al., 1981)
- Size is flexible (about 50 – 90 cm)
- Comfortable **interpersonal distance (IPD)** is a compromise of approach and avoidance forces (Argyle & Dean, 1965).
- PPS and IPD share common mechanisms of self-protection.
- **COVID-19 pandemic** has impacted distancing behaviors.
- **Face mask** may serve as a contamination or a pro-social signal.
- **Evidence mixed:** wearing a mask **reduces PPS** (Serino et al., 2021) and **IPD** (Calbi et al., 2021; Cartaud et al., 2020) **OR enlarges PPS** (Sakuma & Ikeda, 2021) and **IPD** (Seres et al., 2020, Welsch et al., 2021).

## Materials and methods



- Online Experiment (on Gorilla)
- Stimuli: pictures of two individuals in following conditions (s. Figure 1 for sample stimuli):
  - ☑ Distance: 50 cm, 90 cm, 120 cm
  - ☑ Mask: mask, no mask
  - ☑ Interaction: shaking hands, no interaction
- Task: Press Q (or P, counterbalanced) if at least one of the individuals in the picture is a man (or a woman, counterbalanced) -> RT measured
- 235 participants (18 males, mean age 24 y. o.)




Figure 1. Sample stimuli. Panel a: female-male, 90 cm, interaction, no mask. Panel b: male-male, 50 cm, no interaction, no mask; Panel c: male-female, 150 cm, interaction, mask. Panel d: male-male, 50 cm, interaction, mask.

## Results

### Main effects:

**face mask**,  $F(1, 215) = 39.15, \eta^2 = .15, p < .001$  -> processing advantage for pictures with a face mask (mean  $RT = 628.35, SD = 70.50$ ); **distance**,  $F(2, 422) = 49.11, \eta^2 = .19, p < .001$

- fastest RTs for 90 cm (mean  $RT = 624.75, SD = 70.09$ )
- 50 cm (mean  $RT = 632.68, SD = 69.80$ )
- 150 cm (mean  $RT = 638.60, SD = 69.86$ )

**interaction style**,  $F(1, 215) = 4.53, \eta^2 = .02, p = .03$  -> faster RT for stimuli without interaction (mean  $RT = 630.78$ ,




Figure 2. Reaction times [ms] as a function of face mask and interaction style in all three distances (50 cm/near, 90 cm/middle, 150 cm/far). Panel A: 150 cm. Panel B: 90 cm. Panel C: 50 cm. Error bars denote one standard deviation of the mean.

### Interaction effects:

**face mask\* distance**,  $F(2, 429) = 3.19, \eta^2 = .02, p = .04$ ,

**distance \* interaction**,  $F(2, 424) = 36.10, \eta^2 = .14, p < .001$

**face mask \* distance \* interaction**,  $F(2, 425) = 41.46, \eta^2 = .16, p < .001$ .

-> fastest RT with a face mask and no interaction at 90 cm; slowest RT with face mask, with no interaction and at a distance of 150 cm (see Figure 2),

## Discussion

- ☑ Face masks as a symbol of protection facilitate social behavior.
- ☑ Avoidance behavior when interaction.
- ☑ Individuals are perceived more positively when the IPD is 90 cm.
- ☑ Further factors: general perception of COVID-19 as danger, especially for the elderly, having been infected.

## References

- Argyle, M., & Dean, J. (1965). Eye-contact, distance and affiliation. *Sociometry*, 289-304. <https://doi.org/10.2307/2796977>
- Calbi, M., Langiulli, N., Ferroni, F., Montalti, M., Kolesnikov, A., Gallese, V., & Umiltà, M. A. (2021). The consequences of COVID-19 on social interactions: an online study on face covering. *Scientific reports*, 11(1), 1-10. <https://doi.org/10.1038/s41598-021-01780-w>
- Cartaud, A., Quesque, F., & Coello, Y. (2020). Wearing a face mask against COVID-19 results in a reduction of social distancing. *PLoS one*, 15(12), e0243023. <https://doi.org/10.1371/journal.pone.0243023>
- Rizzolatti, C., Scandolara, C., Matelli, M., & Gentilucci, M. (1981). Afferent properties of pericruciate neurons in macaque monkeys. II. Visual responses. *Behavioural brain research*, 2(2), 147-163. [https://doi.org/10.1016/0165-0270\(81\)90053-x](https://doi.org/10.1016/0165-0270(81)90053-x)
- Sakuma, N., & Ikeda, K. (2021). Effects of modulated emotion and facial masking encounter on personal distance. *Perceptual and Motor Skills*, 128(1), 178-190. <https://doi.org/10.1177/0013164420972495>
- Seres, G., Balleyer, A. H., Cerutti, N., Friedrichsen, J., & Süer, M. (2020). Face mask use and physical distancing before and after mandatory masking: evidence from public waiting lines. SSRN. Preprint posted online, 2. <https://ssrn.com/abstract=3641667>
- Serino, S., Trabanelli, S., Jandus, C., Fellrath, J., Grivaz, P., Paladino, M. P., & Serino, A. (2021). Sharpening of peripersonal space during the COVID-19 pandemic. *Current Biology*. <https://doi.org/10.1016/j.cub.2021.06.004>
- Welsch, R., Wessels, M., Bernhard, C., Thönes, S., & von Castell, C. (2021). Physical distancing and the perception of interpersonal distance in the COVID-19 crisis. *Scientific reports*, 11(1), 1-9. <https://doi.org/10.1038/s41598-021-00761-g>